

largely on starlings and ferrets, experimenting with the latter animals in England as well as in this country.

Not merely the length of exposure to light, but other factors as well, have to do with bird migration and other sex-controlled changes in bird and mammal behavior, Prof. Bissonnette concludes, on the basis of his own and other scientists' researches. The intensity of light is one factor: changes occurred more rapidly under bright electric light than under dim illumination of equal duration. Color is another factor: red stimulates sex-gland development, green retards it, at least in starlings. With ferrets, all visible light is about equally effective. Food is a third: starlings kept

on a strict lenten diet of bran mash developed no interest in mating, though light conditions were made favorable.

Experiments both in England and in America indicate that the light factor becomes effective through the hypophysis, a sex-controlling gland situated just under the brain. The hypophysis is stimulated by light falling on the animal's eye. Blind animals, and animals kept with hooded eyes, are not affected by day-length changes, while "control" specimens with normal vision go through the changes in sex activity in regular predictable fashion, in relation to seasonal variations of lighting.

Science News Letter, November 9, 1935

MARINE BIOLOGY

Starfish Are Dyed Blue To Trace Wanderings

BLUE starfish are staring startled oystermen in the face, along the Long Island Sound oyster beds. The orthodox color for starfish in those waters is pink.

The oystermen, however, are being told what it is all about. (*Fisheries Service Bulletin*, Sept. 3). The starfish have been dyed blue for the same reason that migrating birds have aluminum bands fastened about their legs by scientists: to tell where they came from, and how far they have traveled.

Starfish, as is well known, are among the worst enemies oysters have. They fold themselves over the oysters' shells, smother them until they open up, and then devour them. Starfish devastate oyster beds by the square mile in this way.

During the recent session of Congress, funds for the scientific investigation of starfish and other enemies of oysters were appropriated, and research begun at several points. One of the things Victor Loosanoff, Bureau biologist, wanted to learn was the rate of starfish travel—for starfish can move, despite their not-very-motile appearance.

Hard to Tag

But it proved impossible to fasten the customary types of tags or bands on them. They could wriggle out of any kind of knot, and if a tag were stapled right through an arm the starfish calmly shed the arm and grew another. Starfish can do that kind of thing pretty easily.

But a starfish cannot get outside its own hide so easily, so the scheme was finally hit upon of dyeing them a conspicuous color and then turning them loose in thousands. Nile blue, a powerful anilin dye, proved well adapted for the purpose. One ounce of the dry powder provides enough solution to stain more than 25,000 starfish blue.

The process is very simple. The captured stars are dumped into the dye-vat for about a minute, held on deck for an hour, and pushed back into the sea. It costs far less, in both money and time, than any other method of "tagging" living specimens.

Oystermen and all other persons finding blue starfish are requested to write to the U. S. Bureau of Fisheries Laboratory at Milford, Conn., stating the exact location, depth of water and time of capture.

Science News Letter, November 9, 1935

PUBLIC HEALTH

Nation-Wide Grading Of Milk Urged to Prevent Epidemics

ANATION-WIDE uniform system of grading all milk produced in the country so that consumers would know the quality of the milk they were using, whether at home or traveling, was urged by Leslie C. Frank, sanitary engineer of the U. S. Public Health Service, at the American Public Health Association meeting.

Such a system would be of advantage to both consumer and producer, Mr. Frank pointed out.

There are in this country each year at least 30 to 50 outbreaks of disease resulting from infected milk, Mr. Frank said. Most people who use milk know that not all milk supplies are safe, but they have no way of distinguishing a safe from an unsafe supply.

Milk producers would benefit also from the plan Mr. Frank presented. The problem of the producers is one of over-production, or of production of more milk in certain areas than can be profitably sold. Price-fixing efforts have failed and curtailing production would be unwise, in Mr. Frank's opinion.

From the public health viewpoint, not enough milk is being used in the country as it is.

"The only logical solution of both problems is that consumers and industry members coordinate their efforts in a unified nation-wide program to increase milk consumption to the optimum," Mr. Frank declared. "This can be done best by increasing the desirability and safety of milk.

"The voluntary local adoption and strict enforcement of the uniform Pub-

lic Health Service Milk Ordinance, with the rating of all communities periodically by the State authorities, and with occasional check ratings by the Public Health Service, is offered as the form which the coordinated effort should take. Thus America will not only have safeguarded its most important food-stuff, but will also have promoted the economic welfare of the industry which produces and distributes it."

Science News Letter, November 9, 1935

OCEANOGRAPHY

From the Top Down

● "So far as climbing is concerned, the mountains on the ocean are just the opposite of those on land. Our mountain climbers on the land start out at the foot of the mountain . . . But when a man wishes to explore an ocean mountain he starts at the summit and tries to walk down its sides; for the summit is usually an island like Bermuda."—W. Maxwell Reed in *THE SEA FOR SAM* by W. Maxwell Reed and Wilfred S. Bronson (*Harcourt, Brace & Co.*).

Science News Letter, November 9, 1935

Honey bees are about fifty times as valuable for their work of cross-pollination in orchards and fields as they are for the honey they make.

Making a bonfire of autumn leaves is often short-sighted, warns a Cornell agronomist, for decayed leaves form humus and humus helps soils to hold more of the water they need.