

women who habitually abort are capable of producing living, apparently non-syphilitic children when given specific treatment through each pregnancy.

4. Many more non-syphilitic living children were born when anti-syphilitic treatment was begun before the fifth month of pregnancy than when treatment was delayed.

5. If an early syphilis appears late in pregnancy, some treatment begun at

this period and continued until termination of the pregnancy, even though it is only a small amount, will be of value in the production of a living child.

6. Treatment during a preceding pregnancy is insufficient protection for the present pregnancy, even though the syphilitic woman has a negative blood reaction. It is necessary to treat her throughout each pregnancy in order to insure a living non-syphilitic infant.

*Science News Letter, February 15, 1936*

#### WILD LIFE

## Are Duck, Bobwhite, Deer On Way to Extinction?

### Conference Hears That Even Cottontail Rabbit Is Becoming So Scarce That Replacement Is Needed

**E**XTINCTION'S shadow hovers hawk-like over America's remaining wildlife. Will mallard and bobwhite, deer and beaver, vanish as passenger pigeon, great auk, and heath hen have vanished?

No, was the emphatic answer given at the first session of field biologists and conservationists, gathered for the first North American Wildlife Conference, held in Washington. Several score of wildlife scientists and wildlife administrators spoke their minds on special problems, in papers presented at sectional sessions.

It might be thought that the prolific cottontail rabbit would be about the last animal to cause conservationists worry by diminishing numbers, yet such is not the case. Although in some states the value of the cottontails shot annually is greater than that of all other game put together, the supply is running low enough, in some localities, to necessitate the spending of thousands of dollars a year for breeding and replacements of these animals, Dr. R. E. Trippensee, professor of wildlife management at Massachusetts State College, informed the meeting.

Wildlife conservation and replenishment is not entirely a matter for Federal and State agencies, spending millions of dollars upon thousands of square miles of land. Every farmer can run his own conservation program, and many thousands of them are already doing it, on a really scientific basis. Sparing brushy timber patches on creek bottoms and

upland, encouraging shrubs and low trees where gullies need to be checked, planting patches of such game feeding crops as milo maize and leaving them for the birds to harvest, are some of the background elements of such farm-wildlife projects. Game thus encouraged becomes a paying crop in the end, under several different cooperative schemes in force in various states.

L. G. MacNamara told how it is

worked in New Jersey, Aldo Leopold explained the Wisconsin system, F. B. O'Connell spoke of conditions in Nebraska, from the neighbor state of Iowa came Logan Bennett, and A. L. Clark related developments in Connecticut.

Yellow corn, though not a native plant, seems to be the best thing for bobwhite quail under winter conditions in the West, Prof. Paul L. Errington of Iowa State College told his colleagues. Some weed seeds they eat freely and digest, he said, but such things as sumac fruits and wild-rose hips, often thought of as natural quail food, they resort to only when real starvation threatens. These cut their hunger deaths by about fifty per cent.; but there is little real nutriment in them for the birds.

*Science News Letter, February 15, 1936*

#### GENERAL SCIENCE

### "Circus of Science" Will Tour the Country

**A** CARAVAN of 28 motor trucks containing exhibits of science in industry is going on tour throughout the country, Alfred P. Sloan, Jr., president of General Motors Corporation, has announced.

Frankly using the idea behind the Chicago Century of Progress, the "circus of science" will bring to leading



#### SCIENCE JOINS THE CIRCUS

Robert Audet, theatre manager, uses the arts of the stage to bring home the facts of science to the audience attending the "show" of General Motors Parade of Progress.

cities throughout the nation the latest developments and feats of applied science. A great silver-topped tent will

serve as an exhibit hall and supply the circus atmosphere.

*Science News Letter, February 15, 1936*

#### PHYSIOLOGY

## Brain Waves Used in Tracing Activity of Brain Centers

**Clue to Areas of Brain Deterioration in Those With Mental Disease May be Given by Variation in Waves**

**T**HE CHIEF regions of rhythmic activity in the brain can now be located roughly through science's latest researches upon brain waves, the electrical currents that originate in the human brain. Drs. H. H. Jasper and H. L. Andres, of Brown University's psychology department and the Emma Pendleton Bradley Home at Providence, R. I., have also used brain waves as pointers to defective and deteriorated parts of the brain not functioning as they should.

Not any closer to "reading minds" by brain waves, nevertheless science seems likely to use the new knowledge gained in understanding thought processes and in studying mental diseases.

The brain waves are not uniform throughout the whole brain, the investigators report. (*Journal of General Psychology*, January.) Two distinct types of brain waves, known to scientists as alpha waves and beta waves, have been studied. The alpha waves seem to predominate in the lower back part of the head, although under certain conditions some individuals show mostly beta waves in this region. The alpha waves are affected differently by light stimulation than are the beta waves.

In the normal person, alpha waves on the right side of the head are alike in frequency and amplitude to those in a similar region of the left side, although they may be somewhat out of phase, one lagging behind the other. This similarity is absent in a mental patient having one side of the brain diseased. Pictures of the electric waves led off the two sides of the brain of sufferers from brain injury, for example, following lead poisoning, showed a great difference between the activity on the two sides of the brain.

Brain waves of the other type, known as beta waves, are faster than the alpha waves. In any individual they have about twice the frequency and half the amplitude of that person's alpha waves.

They are more predominant in the area at the top of the head known as the Rolandic region. This is the part of the brain governing movement of the body. Occasionally outbursts of alpha waves are also found in this home of the beta waves.

In epileptic patients, the brain wave records during a seizure show what some scientists have likened to a "neurological thunder storm." Drs. Jasper and Andres have confirmed the finding that even when the patient is not undergoing a seizure, his susceptibility to them is betrayed by his brain wave record.

Sporadic seizure waves localized in the motor region of the brain, for example, when no seizure is present produce a similar pattern on the record paper to that seen in all parts of the brain during a generalized convulsion. The method of localized recording described by these authors permits the following of these seizure waves as they spread from one region of the brain to another.

*Science News Letter, February 15, 1936*

#### ARCHAEOLOGY

## Find Tools Believed Left In America During Ice Age

**S**TRONG evidence tending to show that man existed in North America before the end of the Ice Age is announced by Prof. Paul MacClintock, of the Princeton University Geology Department. The evidence is reported to be human implements found in the White River region of South Dakota and Nebraska and deposited there before the time of the last glacial advance.

In the field with Prof. A. L. Lugen of the University of Nebraska and assisted by Justus S. Templeton, a Princeton senior from Dixon, Illinois, Prof. MacClintock discovered last summer varved sediments in the bed of an extinct lake formed when the valley of

the White River was dammed by the last advance of the ice sheet. If these sediments are contemporaneous with the ice sheet, they figured that man-made artifacts found in or below the lake sediments would prove that man was there before the ice.

Accordingly, C. B. Schultz and his party from the Nebraska State Museum, under E. H. Barbour, dug below the sediments and found not only many artifacts, but scores of hearth-pits containing charcoal, burned stones, and burned bones of extinct animals. The pits, which are two or three feet wide and one or two feet deep, seem to have been used to preserve the fire from day to day. The artifacts are of the Yuma type.

While the evidence thus far uncovered is not absolutely conclusive, it is believed that another season will either prove or definitely disprove the hypothesis that man appeared on this continent before the end of the Ice Age. Older theories have leaned strongly to the belief that man did not arrive in North America until the last glaciers had melted sufficiently to allow a crossing from Asia.

*Science News Letter, February 15, 1936*

#### ARCHAEOLOGY

## Cretan Images Confirm Truth of Ancient Writings

**P**EASANTS' discovery of a ruined building near Dreros has led archaeologists to unearth a remarkable temple, believed to be the oldest Greek temple in Crete.

The temple dates from the eighth century B.C., when Crete's own brilliant civilization had faded, and the Greeks were rising to be leaders of culture.

Outstanding among the temple's contents are bronze statues of two goddesses and a naked child god. These religious images were made by hammering thin plates of bronze to shape, possibly around a wooden core, and fastening the bronze in place with pins. Ancient Greek writers described such a process, but never before have archaeologists found actual statues thus hammered. Greek sculptures known today were made by casting.

The temple, measuring 30 by 20 feet, contains a square hearth still covered with ashes, and a stone base on which a wooden column once stood.

*Science News Letter, February 15, 1936*

Butterflies are reported increasingly scarce in British forests.