

MEDICINE

New Approach to Malaria

Tetanus toxoid, given in conjunction with anti-malarial drugs, stimulates the body's defense forces against the disease.

➤ A NEW approach to the world-wide fight against malaria has been discovered by Dr. Eusebio Y. Garcia, senior malariologist of the medical research clinic at Binan in the Philippines.

Bouts of chills and fever did not come back as soon to plague the malaria victim when he was given, along with atabrine or chloroquine, doses of tetanus toxoid, Dr. Garcia found.

Tetanus toxoid is the detoxified poison of the tetanus, or lockjaw, germs. Members of the U. S. armed forces during the war knew it as one of the many "shots" given them for protection against diseases they might encounter. Modern mothers know it is given their babies along with shots against diphtheria and

whooping cough, so that when Junior steps on a nail he will be protected against the lockjaw danger grandmother feared from stepping on nails.

News of Dr. Garcia's work comes to American scientists through the New York Academy of Sciences which has awarded him a \$200 A. Cressy Morrison prize for it.

The use of tetanus toxoid is different from chemical, or drug, treatment of malaria and different, also, from use of germ chemicals, such as penicillin, to fight other germs in the body. Tetanus toxoid, he believes, acts as a stimulator to the body's own defensive forces. Immune mechanisms is the term scientists use for these defensive forces. They are

stimulated and enhanced, he thinks, by the tetanus toxoid, so that relapses in malaria patients after treatment do not come as often.

The significance of Dr. Garcia's work lies in the fact that he is using tetanus germs, which are bacteria, to fight germs which are protozoa and completely out of the bacterial germs' class. It is something like using a featherweight to stop Joe Louis, only more so because the bacteria and the protozoa are as different as plants and animals. Dr. Garcia's results are said to be the first successful attempts to check multiplication of protozoan parasites by stimulating immune mechanisms through a bacterial product.

Science News Letter, December 27, 1947

SCIENCE NEWS LETTER

Vol. 52 DECEMBER 27, 1947 No. 26

The weekly summary of Current Science, published every Saturday by SCIENCE SERVICE, Inc., 1719 N St., N. W., Washington 6, D. C., North 2255. Edited by WATSON DAVIS.

Subscriptions—\$5.00 a year; two years, \$8.00; 15 cents a copy. Back numbers more than six months old, if still available, 25 cents.

Copyright, 1947, by Science Service, Inc. Reproduction of any portion of SCIENCE NEWS LETTER is strictly prohibited. Newspapers, magazines and other publications are invited to avail themselves of the numerous syndicate services issued by Science Service.

Entered as second class matter at the post office at Washington, D. C., under the Act of March 3, 1879. Established in mimeographed form March 18, 1922. Title registered as trademark, U. S. and Canadian Patent Offices. Indexed in Readers' Guide to Periodical Literature, Abridged Guide, and the Engineering Index.

Member Audit Bureau of Circulation. Advertising Representatives: Howland and Howland, Inc., 393 7th Ave., N.Y.C., Pennsylvania 6-5566 and 360 N. Michigan Ave., Chicago, State 4439.

SCIENCE SERVICE

The Institution for the Popularization of Science organized 1921 as a non-profit corporation.

Board of Trustees—Nominated by the American Association for the Advancement of Science: Edwin G. Conklin, American Philosophical Society. **Nominated by the National Academy of Sciences:** Harlow Shapley, Harvard College Observatory; Warren H. Lewis, Wistar Institute; R. A. Millikan, California Institute of Technology. **Nominated by the National Research Council:** Hugh S. Taylor, Princeton University; Ross G. Harrison, Yale University; Alexander Wetmore, Secretary, Smithsonian Institution. **Nominated by the Journalistic Profession:** A. H. Kirchhofer, Buffalo Evening News; Neil H. Swanson, Executive Editor, Sun Papers; O. W. Riegel, Washington and Lee School of Journalism. **Nominated by the E. W. Scripps Estate:** Max B. Cook, Scripps Howard Newspapers; H. L. Smithton, Executive Agent of E. W. Scripps Trust; Frank R. Ford, Evansville Press.

Officers—President: Harlow Shapley, **Vice President and Chairman of Executive Committee:** Alexander Wetmore. **Treasurer:** O. W. Riegel. **Secretary:** Watson Davis.

Staff—Director: Watson Davis. **Writers:** Frank Thone, Jane Stafford, A. C. Monahan, Marjorie Van de Water, Martha G. Morrow, Ron Ross. **Science Clubs of America:** Joseph H. Kraus, Margaret E. Patterson. **Photography:** Fremont Davis. **Sales and Advertising:** Hallie Jenkins. **Production:** Priscilla Howe.

MEDICINE

Painless Shots Possible

By chilling the skin with a new metal applicator, it is possible to block the pain sensation felt by patients while getting injections.

➤ THE needle really won't hurt next time you get a shot of penicillin or some other hypodermically-injected drug if your doctor uses a method devised by three Harvard Medical School surgeons.

The method is to chill the skin briefly to about a freezing temperature before sticking in the hypodermic needle. The chilling is done with a pre-chilled metal applicator of about one-third to one-half inch diameter at its business end. The chilling does not hurt and does block the pain sensation felt by many patients, especially sick children, when the needle is pushed through the skin.

The chilling applicator and its successful use are described by Drs. Franc D. Ingraham, Donald D. Matson and Robert P. Woods in the *New England Journal of Medicine* (Nov. 20).

After trials with carbon dioxide snow, they developed a simple instrument consisting of a brass cylinder about nine inches long and about two inches in diameter. This is closed at one end, and

to the closed end is soldered a smaller brass cylinder. This smaller cylinder which becomes the business end of the instrument, is filled with solder, which is a rapid conductor of heat. The larger cylinder is wrapped with felt and filled with cracked ice to which anhydrous calcium chloride is added. A cork closes it. The contact end of the small cylinder is covered with a closely fitting hollowed-out rubber cap. For use, this rubber cap is removed, the end is wiped clean with alcohol or other antiseptic, and then applied for about 45 to 60 seconds to the previously cleansed skin where the hypodermic injection is to be given.

The instrument can also be used to kill the pain of the needle when blood is taken from a vein in the arm. It is not satisfactory, however, for injections of local anesthetics, presumably because the anesthetic fluid when injected stimulates different nerve endings in the skin from those made insensitive to pain by the chilling.

Science News Letter, December 27, 1947