

PHYSICS

Which of Three H-Bombs?

No information is available concerning the type of "fusion" bomb developed by the U. S. or by Russia. Several kinds are possible.

► ASSUMING THAT Russia does have the hydrogen superbomb (and we must credit Soviet science with the capability), the question is what kind of a hydrogen bomb do they have? What kind do we have?

For there are several possible "fusion" bombs that operate by what is called "thermonuclear reactions."

They are all hydrogen bombs, but there are three kinds of hydrogen: 1. The ordinary and plentiful kind. 2. Deuterium, the double weight kind, rare but occurring in nature. 3. Tritium, the triple weight kind that is radioactive and has to be manufactured.

From highly secret tests at Eniwetok, our scientists probably know now which kinds can be used for superbombs. For there is little doubt that we have exploded one or more hydrogen bombs.

Most of those engaging in more or less informed guessing believe that tritium is the superbomb material of choice. Two atoms of tritium (H^3) brought together with sufficient violence would fuse into and form a helium atom (He^4) and give off two neutrons. Neutrons are the uncharged atomic particles that trigger the atom bomb.

There would be a terrific release of energy due to the fact that mass of the two tritiums is slightly more than that of the helium and the neutrons, and that this mass is converted into energy.

But two deuterium atoms similarly thrown together should also fuse, giving in one reaction a tritium atom and an ordinary hydrogen atom, and in an alternate reaction a mass three helium atom and a neutron. Both would release large energy.

There could also be reactions between atoms of deuterium and tritium, which should be violently explosive. Participation of ordinary hydrogen would be likely to slow things down too much.

All the hydrogen bombs presumably would need extremely sudden and high temperatures to be kicked off. A fission bomb of uranium 235 or plutonium (the conventional atomic bomb if anything atomic can be called conventional) is used to trigger a hydrogen bomb. It creates sun-like temperatures, a million degrees or so.

The speed of interaction of the hydrogen material is all important. For the violence of the explosion will throw the material apart and it must do its combining in a few millionths of a second.

The hydrogen bomb is cheaper and more practical from a production standpoint if its material is deuterium, or double-weight hydrogen. This isotope of hydrogen is

separated from water, in which a small amount occurs. The process is somewhat like the way in which the fissionable uranium, isotope 235, is separated from the most plentiful uranium 238 with which it occurs in the natural deposits. Deuterium can be obtained without use of materials that enter into the production of fission or ordinary A-bombs.

But the production of tritium is a drain on the uranium supply. It must be manufactured, probably from lithium metal, by bombardment with neutrons. The fissioning of uranium, or plutonium made from uranium, is the only practical source of neutrons for this purpose. This would be done in an atomic reactor of the same sort that is used to make plutonium, by bombarding uranium 238 with the neutrons from a slowed-down reaction such as occurs in the atomic fission bomb.

Tritium for hydrogen bombs can be made only at the sacrifice of atomic bomb production and a large sacrifice of total explosive power. It is necessary to forego some of the total punching power to obtain the super-punch of a bomb that is perhaps a thousand times that of the Hiroshima bomb,

a superbomb capable of devastating vast territory.


If deuterium can be made to explode, hydrogen bomb production is not dependent upon the uranium supply except to provide the triggering fission bombs.

Tritium is radioactive and half of its disintegrates in about 11 years. Making it and stockpiling for use in the distant future is wasteful because of this natural disintegration.

All of the hydrogens as elements are gases which are unhandy to use, but they can be combined with other elements into solids. Plutonium tritide or plutonium deuteride would give both the fissionable and the fusionable atomic energy elements in the same compound. Perhaps they are packaged in this way for the hydrogen bomb.

A hydrogen bomb would probably be detectable by the large amount of radioactive carbon created in the gigantic explosion. The nitrogen in the atmosphere bombarded by neutrons given off by the superbomb would be turned into carbon 14, just as the cosmic rays create continuously this kind of carbon in very small amounts in the upper atmosphere. This radioactive material comes down to near the earth's surface and can be used to date things that contain carbon.

If a hundred or so superbombs were exploded, they might so poison the atmosphere with radioactivity as to affect human life, increasing the cancer incidence. If cobalt were added to the hydrogen bomb with fiendish intent, the radioactive cobalt-60 produced and spread in the air would



NAVY'S SEA-DART—The first combat-type aircraft to use water skis for improved rough water take-off and landing operations is the Navy's XF2Y-1 Sea-Dart, experimental delta-wing jet fighter shown here. Retracted, the hydro-skis disappear into the hull to give the plane a missile-like look. (See SNL, Jan. 3, p. 6.)

have the effect of so much deadly radium. Life itself on earth would be jeopardized.

When and if the government tells what has been discovered by the superbomb research and development, we shall be able to form a better judgment as to the hazards of superbombs.

Defense of the civilian population is little changed by a merely bigger bomb. As the bombs become more powerful, the possibility of making peace and keeping out of war with the Soviets becomes more important.

Science News Letter, August 22, 1953

CYTOLOGY

New Way to Study Cell-Virus Battle

► THE BATTLE between body cells and invading disease virus can, it is hoped, be watched and recorded in moving pictures by a new method devised by Drs. Ralph Buchsbaum and Kenneth Wertman and graduate students Evelyn Dwyer, Ronald Gillette and Brother Joseph Kuntz at the University of Pittsburgh.

Body cells are kept alive under a microscope inside a plastic incubator. The incubator keeps the cells at the temperature needed for life and growth, and nourishment is given through special fluids that flow under suitable controls over the cells.

By changing the composition of the fluid or by adding to it a specific virus, the scientists hope to be able to see under the microscope and record with their movie camera the reactions of the cells to changes in nourishment or under attack by disease viruses.

At present chick embryo tissue is under study. Later the scientists hope to study human tissue cells. The project has financial support from the Office of Naval Research, the Atomic Energy Commission and the Army Chemical Center.

Dr. Buchsbaum will go to Rome, Italy, in September, to report on the project before a meeting of the International Congress of Micro-Biology.

Science News Letter, August 22, 1953

PHARMACOLOGY

New Drug Stops Coughs

► GOOD RESULTS with a new synthetic drug that stops coughing spasms without making addicts of its users are reported by Drs. Leo J. Cass and Willem S. Frederik of Harvard Medical School in the *New England Journal of Medicine* (Aug. 12).

The new drug is known chemically as dextromethorphan hydrobromide. (See SNL, Jan. 3, p. 7.)

The drug was made and is still in the process of development and investigation by scientists in the Nutley, N. J., laboratories of Hoffmann-La Roche, Inc. Chemically, it is the dextro isomer of the powerful synthetic pain-killer, Dromoran. But it lacks the pain-relieving ability of Dromoran as well as addiction-causing feature.

PUBLIC HEALTH

Alert on TB Meningitis

► A WARNING to parents to be alert to symptoms of tuberculous meningitis in their children is given by the National Tuberculosis Association. In this kind of tuberculosis, the TB germs attack the meninges, which are the membranes covering the brain and spinal cord.

Before the discovery of streptomycin, this form of TB was practically 100% fatal. Even with this drug, the disease is often fatal and always serious. As in other forms of tuberculosis, early treatment is important to saving the patient's life.

Young children who get tuberculosis from a grown person frequently develop the meningitis form of the disease. For that reason, parents should be particularly alert to early symptoms of the disease if their children have been in contact with a grown-up who has tuberculosis. Grown persons also may develop this form of TB, and it may be a complication of pulmonary tuberculosis, spreading to the meninges from the lung.

The early symptoms of tuberculous meningitis do not seem particularly significant and may not arouse suspicion. The child may be irritable, have a low-grade fever, and may have difficulty in feeding, with vomiting on occasion. Later, the child may slip into a comatose (unconscious) or semi-comatose state. The child who has convulsions may be fortunate, for the family is shocked into action and a doctor is consulted.

The first person to notice such symptoms in the child should be aware of the possibility of tuberculous meningitis.

About 75% of the time, one specialist says, the adult contact from whom the child caught tuberculosis was in the immediate family. When there is a question about tuberculous meningitis in a child but the diagnosis is not immediately certain, members of the family should be X-rayed. Exam-

ination of the family may not only uncover tuberculosis in the adult but aid in the diagnosis of the child's ailment.

One child brought to this specialist had a father with only a slight "cigarette" cough. But when he was X-rayed a huge cavity was found in his lung. His wife had tuberculous pneumonia. Neither of them had seen a physician. The child died with tuberculous meningitis within two weeks.

Science News Letter, August 22, 1953

SCIENCE NEWS LETTER

VOL. 64 AUGUST 22, 1953 NO. 8

The Weekly Summary of Current Science, published every Saturday by SCIENCE SERVICE, Inc., 1719 N St., N. W., Washington 6, D. C., North 7-2255. Edited by WATSON DAVIS.

Subscription rates: 1 yr., \$5.50; 2 yrs., \$10.00; 3 yrs., \$14.50; single copy, 15 cents, more than six months old, 25 cents. No charge for foreign postage.

Change of address: Three weeks notice is required. When ordering a change please state exactly how magazine is now addressed. Your new address should include postal zone number if you have one.

Copyright, 1953, by Science Service, Inc. Reproduction of any portion of SCIENCE NEWS LETTER is strictly prohibited. Newspapers, magazines and other publications are invited to avail themselves of the numerous syndicate services issued by Science Service. Science Service also publishes CHEMISTRY (monthly) and THINGS OF Science (monthly).

Printed in U. S. A. Entered as second class matter at the post office at Washington, D. C., under the act of March 3, 1879. Acceptance for mailing at the special rate of postage provided for by Sec. 34.40, P. L. and R., 1948 Edition, paragraph (d) (act of February 28, 1925; 39 U. S. Code 283), authorized February 28, 1950. Established in mimeographed form March 18, 1922. Title registered as trademark, U. S. and Canadian Patent Offices. Indexed in Readers' Guide to Periodical Literature, Abridged Guide, and the Engineering Index.



Member Audit Bureau of Circulation. Advertising Representatives: Howland and Howland, Inc., 1 E. 54th St., New York 22, ELdorado 5-5666, and 360 N. Michigan Ave., Chicago 11, State 2-4822.

SCIENCE SERVICE

The Institution for the Popularization of Science organized 1921 as a non-profit corporation.

Board of Trustees—Nominated by the American Association for the Advancement of Science: Ferdinandus Payne, National Science Foundation; Karl Lark-Harowitz, Purdue University; Kirtley F. Mather, Harvard University. Nominated by the National Academy of Sciences: Harlow Shapley, Harvard College Observatory; R. A. Millikan, California Institute of Technology; Homer W. Smith, New York University. Nominated by the National Research Council: Leonard Carmichael, Smithsonian Institution; Ross G. Harrison, Yale University; Duane Roller, Hughes Aircraft Co. Nominated by the Journalistic Profession: A. H. Kirchofer, Buffalo Evening News; Neil H. Swanson, Baltimore Sun Papers; O. W. Riegel, Lee Memorial Journalism Foundation. Nominated by the E. W. Scripps Estate: John T. O'Rourke, Washington Daily News; Charles E. Scripps, E. W. Scripps Trust; Edward J. Meeman, Memphis Press-Scimitar.

Officers—President: Harlow Shapley; Vice President and Chairman of Executive Committee: Leonard Carmichael; Treasurer: O. W. Riegel; Secretary: Watson Davis.

Staff—Director: Watson Davis. Writers: Jane Stafford, Marjorie Van de Water, Ann Ewing, Allen Long, Horace Loftin. Science Clubs of America: Joseph H. Kraus, Margaret E. Patterson. Photography: Fremont Davis. Sales and Advertising: Hallie Jenkins. Production: Priscilla Howe. Interlingua Division in New York: Alexander Gode, Hugh E. Blair, 80 E. 11th St., GRameray 3-5410.

Science News Letter, August 22, 1953