

## PSYCHOLOGY

# Pick Future Delinquents

► TESTS THAT show promise of spotting potential juvenile delinquents before they become delinquent are now being developed at the University of California.

Dr. T. R. Sarbin, associate professor of psychology, is working to develop a battery of special tests to pick out potential delinquents before they reach high school age. If this can be done, such youngsters can be given special attention to prevent them from becoming social problems.

For the past two years the psychologist has been conducting studies of 100 boys between the ages of 14 and 17. About half are normal boys from a city high school; the remainder are psychopathic delinquents from state institutions.

The object is to develop simple tests that show significant differences between normals and delinquents. Then when youngsters give answers characteristic of the abnormal patterns, they may be potential delinquents.

Two tests are already promising. In one, the youngster copies geometrical figures on a piece of paper. When no restriction is placed on the copying, delinquents respond as accurately as normal boys. But when

limits are imposed, such as drawing the figures on small pieces of paper, the accuracy of the delinquents breaks down immediately. This indicates the inability of delinquents to tolerate limits on their behavior.

In a second test, youngsters see motion pictures in which other youngsters make decisions about situations they are faced with. When the picture is over, the subject is told that the boy in the picture was given a preference test concerning occupations and recreational interests. The subject is then asked to guess what choices the boy in the movie made. The subject is then told whether he is right or wrong about each guess, and is given additional items in the preference test.

The psychologists find that psychopathic delinquents do not seem to "learn" how to make correct analyses of others, but that normal youngsters make better guesses as they are given more experience.

Collaborators with Dr. Sarbin are Dr. Norman Livson, Bela Baker and D. S. Jones of the University of California, and Prof. Paul Wallin, Stanford sociologist.

Science News Letter, January 2, 1954

damage is increased by spray applications from airplanes.

The situation has caused cotton states to take elaborate measures to protect cotton growers. In some states purchases of 2,4-D, 2,4,5-T and related hormone-type herbicides must be registered as in the case of narcotic drugs.

Texas requires that a surety bond of \$20,000 be posted before permission may be obtained to apply 2,4-D. Other regulations specify minimum distances from susceptible crops.

Science News Letter, January 2, 1954

## SCIENCE NEWS LETTER

VOL. 65 JANUARY 2, 1954 NO. 1

The Weekly Summary of Current Science, published every Saturday by SCIENCE SERVICE, Inc., 1719 N. St., N. W., Washington 6, D. C., NORTH 7-2255. Edited by WATSON DAVIS.

Subscription rates: 1 yr., \$5.50; 2 yrs., \$10.00; 3 yrs., \$14.50; single copy, 15 cents, more than six months old, 25 cents. No charge for foreign postage.

Change of address: Three weeks notice is required. When ordering a change please state exactly how magazine is now addressed. Your new address should include postal zone number if you have one.

Copyright, 1953, by Science Service, Inc. Reproduction of any portion of SCIENCE NEWS LETTER is strictly prohibited. Newspapers, magazines and other publications are invited to avail themselves of the numerous syndicate services issued by Science Service. Science Service also publishes CHEMISTRY (monthly) and THINGS of Science (monthly).

Printed in U. S. A. Entered as second class matter at the post office at Washington, D. C., under the act of March 3, 1879. Acceptance for mailing at the special rate of postage provided for by Sec. 34.40, P. L. and R., 1948 Edition, paragraph (d) (act of February 28, 1925; 39 U. S. Code 283), authorized February 28, 1950. Established in mimeographed form March 18, 1922. Title registered as trademark, U. S. and Canadian Patent Offices. Indexed in Readers' Guide to Periodical Literature, Abridged Guide, and the Engineering Index.



Member Audit Bureau of Circulation. Advertising Representatives: Howland and Howland, Inc., 1 E. 54th St., New York 22, Eldorado 5-5666, and 360 N. Michigan Ave., Chicago 11, State 2-4822.

### SCIENCE SERVICE

The Institution for the Popularization of Science organized 1921 as a non-profit corporation.

Board of Trustees—Nominated by the American Association for the Advancement of Science: Ferdinandus Payne, National Science Foundation; Karl Lark-Horowitz, Purdue University; Kirtley F. Mather, Harvard University. Nominated by the National Academy of Sciences; Harlow Shapley, Harvard College Observatory; Homer W. Smith, New York University. Nominated by the National Research Council: Leonard Carmichael, Smithsonian Institution; Ross G. Harrison, Yale University; Duane Roller, Hughes Aircraft Co. Nominated by the Journalistic Profession: A. H. Kirchofer, Buffalo Evening News; Neil H. Swanson, Baltimore Sun Papers; O. W. Riegel, Lee Memorial Journalism Foundation. Nominated by the E. W. Scripps Estate: John T. O'Rourke, Washington Daily News; Charles E. Scripps, E. W. Scripps Trust; Edward J. Meeman, Memphis Press-Scimitar.

Officers—President: Harlow Shapley; Vice President and Chairman of Executive Committee: Leonard Carmichael; Treasurer: O. W. Riegel; Secretary: Watson Davis.

Staff—Director: Watson Davis. Writers: Jane Stafford, Marjorie Van de Water, Ann Ewing, Allen Long, Clare Cotton. Science Clubs of America: Joseph H. Kraus, Margaret E. Patterson. Photography: Fremont Davis. Sales and Advertising: Hallie Jenkins. Production: Priscilla Howe. Interlingua Division in New York: Alexander Gode, Hugh E. Blair, 80 E. 11th St., GRamercy 3-5410.

## ENTOMOLOGY

# Pink Bollworm Spreads

► FRANTIC EFFORTS by cotton-belt states have slowed but not halted the spread of destructive pink bollworms. The U. S. Department of Agriculture's Bureau of Entomology and Plant Quarantine has reported discovery of the insect in Arkansas for the first time.

The bollworm is rated as potentially equal to the billion-dollar boll weevil in destructiveness. In 1952, it did more than \$28,000,000 damage in 38 south Texas counties.

Pink bollworm is now largely confined to parts of Texas, Oklahoma, Louisiana and Arizona. Each year, however, its range of destruction is slightly increased.

Unusual weather conditions have contributed to the recent quickened spread in the opinion of experts. These conditions have held back control measures at critical periods of the bollworm's life cycle, allowing a large increase in the bollworm population.

States and counties have drastic quarantine regulations to control the insect. Border patrols this year intercepted several hundred specimens being transported out of infested areas in cotton seeds and plants.

Pink bollworm was first found in Hearn, Tex., in 1917. Since that time its spread has been slow but sure. In addition to spreading by transportation of infected cotton, the adult moths have a small flying

range that may be greatly extended if they are caught in strong air currents.

Every known method of suppression, from insecticides to natural parasites, is being used in the fight against the bollworm. Compared to the quick spread of the boll weevil, the bollworm has been held down for many years, but the fight is far from over for hard-pressed cotton growers.

Science News Letter, January 2, 1954

## AGRICULTURE

## Weed Killer Causes Cotton-Rice War

► WAR BETWEEN rice and cotton growers in the delta area of Mississippi has broken out over the weedkiller, 2,4-D, Dr. Ross E. Hutchins of Mississippi State College reports in *Science* (Dec. 25, 1953).

Rice growers have been using the herbicide to control coffee-weed in their fields. The difficulty is that cotton is "almost fantastically sensitive to 2,4-D and related compounds," Dr. Hutchins says.

Cotton growers now have damage suits totaling several hundred thousand dollars before the courts.

Dr. Hutchins points out that it is not unusual for severe damage to cotton plants to occur four or five miles away from ground applications of 2,4-D, and that this