

their genetic information (RNA) into DNA? Then might the DNA copies of the viral genes be inserted into the DNA of host cells? And might the viral DNA make the cells cancerous?

But research compiled during the past year or two somewhat deflates that hope, or at least makes it harder to document. This fact was brought home last week at a meeting of the American Chemical Society in Chicago.

The reverse transcriptase enzyme, which appeared to be a unique product of RNA tumor viruses, has now been found in noncancer viruses as well. It has also been found in noncancerous cells. It now becomes clear, says Temin, that the enzyme does not always correlate with cancer.

In fact, even RNA tumor viruses may not necessarily lead to cancer. DNA copies of viral RNA are now known to incorporate themselves into the DNA of host cells. But even when this happens, cells do not necessarily become cancerous. "Integration of viral-specific DNA in itself is not the cause of cancer," reports Harold E. Varmus of the University of California at San Francisco.

Have investigators given up hope, then, that RNA-directed DNA synthesis might provide a key to the cancer process? Not really. RNA virus-produced reverse transcriptase does not always turn cells into cancer cells. But transformation of cells by RNA tumor viruses is always accompanied—as far as P. K. Vogt of the University of California at Los Angeles can tell—by viral DNA. The viral DNA has presumably been made by the reverse transcriptase enzyme. So the reverse transcriptase enzyme is necessary, if not sufficient, to cause cancer.

The implication of reverse transcriptase in the cancer process, then, has been amply demonstrated. But what the implication means will only become fully evident when scientists get at the basic genes of viral RNA and identify which genes might transform cells into cancer cells. □



Joan Arehart-Treichel

Temin: It's not a cancer exclusive.

september 8, 1973

Mesons on the mesa: First from the factory

About 10 years ago physicists began to seek the construction of the so-called meson factories. These are machines that would accelerate beams of protons and drive them against targets so as to produce copious beams of mesons, particularly pi mesons. Pi mesons are useful as probes of the atomic nucleus, and nuclear physics stands to benefit greatly from meson factories. So would medicine: Pi mesons have a potential use in the destruction of certain kinds of tumors.

The Clinton B. Anderson Los Alamos Meson Physics Facility (LAMPF) located on top of a mesa at Los Alamos, N.M., is the one project to provide such a meson factory in the United States. Last week it produced its first pi mesons. The event climaxed five years of design, engineering and construction.

The pi mesons were produced in beam area "A," one of three beam areas at the end of the protons' 1,800-foot flight. The area had been declared ready on Aug. 25 and received its first proton beam, of 447 million electronvolts (MeV) energy on Aug. 26. The first pi mesons were produced on Aug. 27. The ultimate plan is to use protons of 800 MeV energy.

Participants in the meson production include Don Cochran, Robert Macek, Robert Burman, Don Hagerman and Mahlon Wilson of Los Alamos Scientific Laboratory and Mark Jakobson of the University of Montana, chairman of the LAMPF Users Group.

The first pi mesons were the signal to begin an effort that includes tuning the main and secondary beam lines and preparing the secondary lines, where the mesons will appear, for experiments. It is expected to take several months. □


Are the outer planets 'failed' stars?

The planets of the solar system can be divided broadly into two classes: the terrestrial (Mercury, Venus, Earth and Mars) and the Jovian (Jupiter, Saturn, Uranus and Neptune). (Pluto is left out because too little is known about it.) The terrestrial planets are small, compact bodies with densities greater than about 4 grams per cubic centimeter. The Jovians are large and diffuse with densities less than 2.25 grams per cubic centimeter. The terrestrials all lie within 1.52 astronomical units of the sun. (One astronomical unit is the radius of the earth's orbit.) The Jovians only begin after a sizable gap: Jupiter, the nearest Jovian to the sun, is 5.20



LASL

LAMPF, meson factory on the mesa.



LASL

Inside the tunnel ready for the beam.

astronomical units away from the sun.

These differences have excited much speculation among astronomers, and the suggestion has been made several times that Jupiter looks more like a small star than a planet. Now comes D. McNally of the University of London Observatory, who argues that all four of the Jovians are "failed" stars. Specifically, he suggests "that the terrestrial planets formed in a high-density shell surrounding the proto-Sun and that the Jovian planets are the remnants of other attempts to form stars contemporaneously with the Sun."

If Jupiter, Saturn, Uranus and Neptune are all indeed failed stars, they failed almost at the point of becoming stars. What inhibited their final collapse into the stellar state? Clearly it was not proximity to the sun: Binary stars are