

member John Oro of the University of Houston, perhaps tightening the known constraints on planets where life as we know it can form. Gravity, he says, will have to be sufficient to retain the volatile elements; atmospheres may be required to possess enough ozone to keep out ultraviolet radiation, and there may have to be

a temperature range—perhaps 0 to 100°C—that is both cool enough to retain water but warm enough to allow the proper chemical reaction rates as pre-life compounds try to get a foothold. “It may sound defeatist,” says one Viking team member, “but the lack of life on Mars would have its interesting side too.” □

Earth's hot past: Delay to evolution?

The farther we go back in time, the more fragmentary is our record of past temperatures on earth. For the oldest geological periods the record is practically nonexistent. But now isotopic analysis of 66 samples of chert from the western and central United States and abroad has added valuable new data to the record, showing that 3 billion years ago the earth was apparently much warmer than it is today, the average temperature perhaps as high as 160°F. The new data indicate that by 1.2 billion years ago, the climatic temperature had declined to 90°F.

The new evidence was obtained at the California Institute of Technology by geochemists Samuel Epstein and L. Paul Knauth. Knauth, then a graduate student, is now on the faculty of Louisiana State University. The findings are reported in the current *GEOCHIMICA ET COSMOCHEMICA ACTA*, the journal of the Geochemical Society and the Meteoritical Society, and in an announcement from Caltech.

Epstein believes the hot ancient temperatures may help explain why advanced forms of life did not evolve on earth until about the latest eon of the planet's existence. “If our estimated ancient temperatures are even nearly correct,” he says, “these temperatures may have been a reason why multicelled organisms did not appear on earth until about one billion years ago. The earth may simply have been too hot for sophisticated life to have evolved.”

It wouldn't have been too warm for bacteria, however, which have been around for at least 3.3 billion years, or the primitive blue-green algae, which followed bacteria on earth.

The Epstein-Knauth data also contain a temperature record of the earth's most recent one billion years. They indicate that the annual temperatures dropped from about 93° to 68°F during the Paleozoic era (600 to 225 million years ago), then climbed to 95° to 104° in the Triassic (225 to 190 million years ago), and declined through the Mesozoic and Cenozoic to values of about 63°. The figures for more recent times correspond to trends, if not actual degrees, shown in previous calculations based on paleontological and other data. The cause of the hot temperatures in earlier times is not known. One suggestion is that the sun was hotter then.

The findings are based on measurements of the relative abundances in the chert of isotopes of oxygen and of hydro-

gen. The isotopic ratios vary according to the temperature of the atmosphere at the time the cherts are formed.

The samples were taken from 11 states, plus southeastern Canada, South Africa and England. The readings turned out to be consistent for geological periods, no matter how far apart the samples were located.

Epstein points out that there are uncertainties in any paleoclimatological method as one goes back in time and says that the results have to be considered tentative.

“However,” he says, “of all the possible interpretations, only those involving changes with time of climatic temperatures come close to accounting fully for the observations.” □

Element 107: U.S. group skeptical

Among scientists interested in the discovery or synthesis of new chemical elements the name of Academician Georgi Flerov is often heard. Flerov and his colleagues in the Joint Institute for Nuclear Studies at Dubna in Russia have specialized in this sort of work for years, and they have outstanding claims to the discovery of several of the elements with atomic numbers greater than 100. Their latest claim is to element 107 and appears in the Russian language express journal *Pis'ma v Redaktsyu ZhETPh*, vol. 23 p. 206. (This will eventually be translated into English as *JETP Letters*.) Flerov and colleagues say they have made a nucleus with 107 protons and 154 neutrons.

Counterclaims to the synthesis of elements 104, 105 and 106 have been entered by an American group that has specialized in synthesizing new elements for decades and has been led by Glenn T. Seaborg and Albert Ghiorso of the Lawrence Berkeley Laboratory. No one, at the moment, makes a counterclaim to discovery of element 107, but Seaborg is extremely skeptical of Flerov's claim. “There isn't sufficient information to assign an atomic number,” he told *SCIENCE NEWS*.

Seaborg's objection is to Flerov's use of spontaneous fission of the supposed new nucleus as evidence for its existence and characteristics. Flerov's experiment began with the bombardment of bismuth nuclei with chromium nuclei in the hope that they would fuse to form the new

element. At first a spontaneously dividing nucleus with a lifetime of 5 seconds was found. This is much longer than theory would expect for the lifetime of 107. Pushing down the response time of the experiment, Flerov's group then found a spontaneously dividing nucleus of about 2 milliseconds lifetime, which comes much closer to the theoretically expected duration of element 107. Further checks led them to conclude that the 2-millisecond nucleus was 107, while the 5-second activity is attributed to nuclei of element 105 produced by alpha decay of element 107.

Seaborg says that this spontaneous fission evidence puts the present claim in the same category as Flerov's claims to several previous heavy elements, that is, highly doubtful. In fact, because of the number of recent claims to discovery of heavy elements (114, 115, 116, 124 and 126) from various sources and the expectation that the spread of heavy-ion accelerators will lead to many more, Seaborg and other persons prominent in the specialty published a manifesto in the Sept. 24 *SCIENCE*, in which they call for the establishment of criteria that a claim to a new element must satisfy before being admitted for consideration. Other signatories are Bernard G. Harvey of LBL, Günter Herrmann of the University of Mainz, Richard W. Hoff of the Lawrence Livermore Laboratory, Darleane C. Hoffman of the Los Alamos Scientific Laboratory, Earl K. Hyde of LBL, Joseph J. Katz of Argonne National Laboratory, O. Lewin Keller Jr. of Oak Ridge National Laboratory and Marc Lefort of the Institut Physique Nucléaire at Orsay, France.

The main point of their argument is that “detection of a spontaneous fission activity and measurement of its half-life cannot per se establish that an element with a new atomic number has been produced.” Chemical tests establishing the candidate's position in the periodic table would be ideal. But this group realizes that chemical procedures with small samples are difficult and so says that if the new element is observed by spontaneous fission or alpha decay or both, “The chemical identification can be confined to separation from all known elements with atomic number greater than lead.” Other acceptable evidence would be observation of characteristic X-rays emitted in connection with the decay of the new element or the placement of the new element in an alpha-particle decay chain with previously known decay products. As a final shot, the manifesto points out that heavy-ion bombardment experiments produce two new kinds of short-lived states, compound nuclei and nuclear molecular systems. These are not really new elements, and to guard against claiming them as such, they urge that composite nuclear systems lasting less than 10^{-14} seconds should not be considered new elements. □