

SCIENCE NEWS®

The Weekly Newsmagazine of Science

A Science Service Publication
Volume 128, No. 21, November 23, 1985

E. G. Sherburne Jr.	Publisher
Joel Greenberg	Editor
Dietrick E. Thomsen	Senior Editor/ Physical Sciences
Laurie Jackson Phillips	Managing Editor
Wendy McCarren	Production/Design Director
Bruce Bower	Behavioral Sciences
Joanne Silberner	Biomedicine
Stefi Weisburd	Earth Sciences
Julie Ann Miller	Life Sciences
Janet Raloff, Ivars Peterson	Policy/Technology
Jonathan Eberhart	Space Sciences
Susan Welch Gilday	Assistant to the Editor
Diane D. Edwards	Science Writer Intern
Jane M. Livermore	Books
Donald R. Harless	Advertising/Business Manager

Copyright © 1985 by Science Service, Inc.,
Editorial and Business Offices,
1719 N St., N.W., Washington, D. C. 20036.
Republication of any portion of SCIENCE NEWS
without written permission of the publisher is
prohibited.

Subscription Department
231 West Center Street, Marion, Ohio 43305

Subscription rate: 1 yr., \$29.50; 2 yrs., \$50.00.
(Foreign postage \$5.00 additional per year.) Change
of address: Four to six weeks' notice is required.
Please state exactly how magazine is to be
addressed. Include zip code. For new subscriptions
only call (1) 800-247-2160. Printed in U.S.A. Second
class postage paid at Washington, D. C. Title
registered as trademark U. S. and Canadian Patent
Offices. Published every Saturday by SCIENCE
SERVICE, Inc. 1719 N St., N.W., Washington, D. C.
20036. (202-785-2255)
ISSN 0036-8423

Letters

Radiation's deadly dose

"New estimates of radiation lethality..." (SN: 10/12/85, p. 232) uses the "old" unit of absorbed dose, the rad. The SI unit, the gray (Gy), should be used starting this year.

The article states that the lethal human radiation dose (LD-50) was previously believed to be 6 Gy (600 rads) to bone marrow. Other sources available to me indicate the value is about 4 Gy (400 rads). This would make a proposed reduction to 1.54 Gy (154 rads) about a factor of 2.5 rather than a factor of 4.

The LD-50 value is for radiation by itself, such as that due to distant fallout from nuclear weapons, with no subsequent medical care. One of the effects of large radiation doses is to reduce the blood's ability to clot and to fight infections. Thus a large radiation dose in conjunction with lacerations is far more serious than either the radiation dose or the laceration by itself.

Most of the victims of the 1945 bombings of

This Week

- 324 Gene-Splice Approvals on Two Fronts
- 324 Single-photon interference seen
- 325 Molecular carbon: Playing 'buckyball'
- 325 AIDS virus: Infection up?
- 326 Colombia volcano: What next?
- 326 Two more sun-grazing comets discovered
- 326 Laskers awarded
- 327 Coffee and the job: Heartfelt insults?
- 327 Lasers powering through coronary arteries

Research Notes

- 328 Biology
- 328 Biomedicine
- 329 Earth Sciences
- 329 Physical Sciences
- 330 Science & Society

Articles

332 Sick Chips

Cover: Two metal leads emanate from the active portion of a failed MOS FET, or metal-oxide semiconductor field-effect transistor. This MOS FET, shown magnified 540 times, was to have been used in the Infrared Astronomy Satellite. In laboratory tests, however, these chips kept failing — short circuiting — from defects caused by electrostatic discharge. The chip shown has four such defects. The most serious, and the one that caused the chip to fail, is the circular hole in the band of metal circuitry about an inch to the right of the upper lead connection. Today's state-of-the-art chips are especially vulnerable to electrical damage that can inflict symptomless yet severely life-shortening damage. (Photo: JPL/Caltech)



Departments

323 Letters

Science Service Institution for the public understanding of science founded 1921; a nonprofit corporation.
Board of Trustees — President, Glenn T. Seaborg; Vice President, Gerald F. Tape; Treasurer, Willis Harlow Shapley; Secretary, Hilleary F. Hoskinson; Joseph W. Berg Jr.; Edward Bliss Jr.; Bowen C. Dees; David A. Goslin; Milton Harris; Elena O. Nightingale; O. W. Riegel; H. Guyford Stever; John Troan; Deborah P. Wolfe.
Director: E. G. Sherburne Jr.; Assistant Director: Dorothy Schriver; Business Manager: Donald R. Harless.

Japan within "892 meters" of ground zero, with radiation doses of 1 to 4 Gy (100 to 400 rads), would have suffered lacerations and/or burns from the heat of the blast (larger doses would have been required to produce serious "radiation burns"). It would seem that the study may have counted deaths from the combined effects of radiation, lacerations and thermal burns, and attributed them to radiation alone.

Carl Fields
Assistant Professor
Nuclear Engineering Technology
Pennsylvania State University
Monaca, Pa.

Regarding the LD-50 cited in the article, Joseph Rotblat notes in his paper: "This value has been deduced partly from animal data and partly from the few radiation accidents, many victims of which have been receiving extensive medical treatment. ... In the U.K. an LD-50 of [6 Gy] to bone marrow — deduced from these accidents — is being used in estimates of radiation casualties in a nuclear war." — J. Raloff

Hypothetical honor

I would like to call into question J.A. Miller's citation of Judah Folkman as the originator of the angiogenesis hypothesis ("Call of the tumor," SN: 10/5/85, p. 213). We believe this honor belongs to Stevan Durovic, M.D., who not only advanced this hypothesis from Argentina during the early 1940s but also supported it with such elegant research that he was sponsored to continue his work under A. C. Ivy, Distinguished Professor of Physiology at University of Illinois and Director of the American Cancer Society, during the next two decades.

Durovic and his colleagues have subsequently been quite successful in developing an organic antiangiogenic agent specific to many tumor cell types, and have demonstrated reversal of tumor growth due to angiocatabolism within 72 hours of its administration.

Charles Wallach
President
Behavioral Research Associates
Canoga Park, Calif.

NOVEMBER 23, 1985

323