

bodies, note Kissileff and his colleagues at the Columbia University College of Physicians and Surgeons.

According to another paper in the same journal, high-protein diets can suppress appetite—at least in monkeys. By a feat of surreptitious plumbing, a Baltimore research team bypassed finicky monkey noses and taste buds to feed four monkeys a dietary supplement containing 50 percent protein. Anticipating that the animals might change their eating habits when presented with the modified diet—and thus throw a primate-wrench into the experiment—the scientists implanted tubes into the monkeys' stomachs and connected them to food-delivery pumps. The monkeys, trained to feed at nozzles connected to other suction-activated pumps, unknowingly triggered both pumps with each oral feeding, thus getting a measured double-dose of food.

The protein-boosted diet caused the monkeys to eat less overall, dropping their total calorie intake by 25 percent, report Judy S. Hannah, Anil K. Dubey and Barbara C. Hansen of the University of Maryland School of Medicine. The finding, they say, adds to the body of evidence suggesting that high-protein diets can aid in weight reduction.

"A high-protein diet does appear to have an appetite-suppressing effect, and of course, if you want to lose weight, that could be beneficial," Hannah says. She warns, however, that people with kidney problems, high blood pressure or diabetes should not undertake a high-protein diet on their own. "In fact, anyone should always consult a physician before making major dietary changes." □

Letters continued from p.131

nary gift for poetic improvisation, which he demonstrated by producing immediate English versions of French, Italian or Latin verse and rhyming, scanning verses on such trivia as children's hats. A world-class procrastinator, he produced some of his best-known essays at the last moment at great speed and apparently never revised his work. Although functioning at a high artistic and intellectual level, Samuel Johnson recalls Dr. Oliver Sacks' "Witty Ticky Ray" in his ability to adapt his disability to creative ends.

Thinking of Johnson as a Touretteur rather than a gross eccentric enhances our appreciation for his achievements and for the 18th-century ability to appreciate talent while overlooking what we today would call pathology.

Donald R. DeGlopper
Bethesda, Md.

Many researchers indeed cite Dr. Johnson as an early documented case of Tourette's syndrome.

— B. Bower

Solar vs. sidereal days

Earth rotates on its axis every 23 hours and 56 minutes, yet an Earth day is 24 hours. Likewise, Earth rotates 366.26 times per year, yet an Earth year has only 365.26 days. The reason for these differences is that, as commonly defined, days are measured relative to the sun ("solar days") while rotations are measured relative to the stars. Consistent with this, if Earth did not rotate at all, it would still experience a day, actually an "inverse" day, for each orbit around the sun ("inverse" meaning the sun would rise in the west and set in the east). This accounts for Earth's "missing" day after 366.26 rotations each year and also for the "extra" 4 minutes (1/366.26 day) required beyond a rotation period in order to complete a day.

For other planets the magnitude of this effect can be fairly astounding—an interesting and

important fact that was not included in your discussion of Mercury's cyclic solar exposure ("Cold message from Mercury's hot poles," SN: 6/16/90, p.375). You state that "a Mercury year lasts but 1½ Mercury days" and that "Mercury [turns] three times on its axis for every two trips it makes around the sun." Clearly, you are referring to "sidereal days" (synonymous with rotations) instead of solar days, despite the discussion's solar context. While sidereal and solar days differ by only 0.27 percent on Earth, they differ by 200 percent on Mercury.

The "inverse" day resulting from Mercury's orbit leaves that planet with a net of only ½ solar day per year. In other words, a stationary observer on Mercury would experience alternating years of daylight and darkness. Thus, each longitude directly faces the sun at only one point (and always at the same point) during each two treks along Mercury's highly elliptical orbit. This results in only two longitudes (180° apart) ever directly facing the sun at the orbit's perihelion, and consequently results in the occurrence of the equatorial "hot poles."

Robert E. Crippen
Jet Propulsion Laboratory
Pasadena, Calif.

CORRECTION

In "Sweet tooth, rotten kid: A theory gone sour" (SN: 8/11/90, p.84), the name of the California State University scientist who commented on the sugar study was misspelled. The correct spelling is Stephen Schoenthaler.

Address communications to:
Editor, SCIENCE NEWS
1719 N St., NW
Washington, D.C. 20036

Please limit letters to 250 words.
All letters subject to editing.

Continued from p.137

burrow D-17 came up speckled with tell-tale white fragments. Sometime late in the '88 season, unbeknownst to the observers, a dark-rumped petrel had laid an egg.

The team repeated the experiment in '89 and '90, adding a total of 140 artificial burrows. Each year, the number of night visitors increased, and more of them lingered through the day. Podolsky and crew found birds sitting in burrows during morning checks in 1989, with burrow D-17 again housing incubating petrels. This summer, the number of nesting petrels tentatively stands at one pair. With the verdict still out on some suspiciously busy burrows, Podolsky is optimistic that the count will rise before the birds head out to sea in a few months for the winter.

At the suggestion that a handful of petrel nests seems a modest return for 220 hand-dug burrows and three seasons of avian advertising blitzes, Podolsky smiles like a salesman

about to close a million-dollar deal. For he knows the nature of his customers.

The dark-rumped petrel often lives into its 30s and takes eight years to mature—an unusually long time for a bird of its size. "When we play our recordings, we don't appear to lure in breeders, we appear to only lure in young prospectors," Podolsky says. "We think they start returning to the island at 2 years old and continue to prospect until they're 8 years old." That leaves a comforting window of time before the return on his investment comes due. And in light of the finding that young petrels have already perused more than 70 percent of the 220 artificial burrows, the prospect of a petrel boomtown becomes easier to envision.

Podolsky and Kress see an expanding global need for their services, and they may be the world's only salesmen to view a rising demand as sad news. Dark-rumped petrels of the Galápagos and Hawaiian islands face a continuing threat from expanding agriculture and introduced predators, they say. And seabird situations around the world look sim-

ilarly grim. Of the approximate 270 species of seabirds worldwide, 30 are listed as endangered or threatened.

"Ultimately, our responsibility as biologists is to maintain the biological richness, the natural heritage of this planet," says Podolsky. "It's like the rivet puller analogy by [ecologist] Paul Ehrlich: You start pulling rivets from an airplane and nothing seems to happen. But there comes a point. . . ."

For now, most of the ornithologists' efforts are mere stopgaps, helping to settle and protect new colonies of endangered seabirds until less predator-infested island homes can be found. For some birds, however, the menaces aren't limited to rats, gulls, pigs and people.

Consider the short-tailed albatross. All but seven of the last 100 or so breeding pairs nest on Tori Shima Island off Japan. "The idea is to lure them to the other side of the island," says Podolsky. In the meantime, the heart of the short-tailed albatross population innocently gathers each fall to lay its precious eggs in the shadow of an active volcano. □