

sure. To validate an assay method, she first withdrew and analyzed blood from 20 healthy men and women.

During each of several visits, the healthy volunteers' blood levels of the marker protein, beta-thromboglobulin (beta-TG), averaged about 30 nanograms per milliliter — except in two of the participants. These individuals showed a more than sevenfold increase in beta-TG on one morning. Upon questioning, each recalled only one unusual thing about the day on which their levels were high: no breakfast.

Intrigued by the possibility of a link between fasting and platelet stickiness, Cifkova initiated a small follow-up study, again using healthy volunteers. Between 10 a.m. and noon on two days no more than one week apart, she and her colleagues assayed beta-TG levels in 19 men and 10 women. Participants ate breakfast before coming in for the first test, but skipped it — maintaining at least a 14-hour fast — before the second test. After initial blood tests on that second day, the volunteers ate a meal. Three hours later, the researchers retested them.

Morning beta-TG levels averaged more than 2½ times higher on the day the group skipped breakfast. After they ate, however, the protein plummeted to levels that “did not significantly differ” from those measured after breakfast on the first day, says Cifkova, who presented her findings last week in Washington, D.C., at the National Conference on Cholesterol and High Blood Pressure Control.

While conceding this study is far from conclusive, she says its results strongly suggest that “overnight fasting and skipping breakfast increases platelet activation and might contribute to the known increased frequency of [heart attacks], sudden death and ischemic stroke during early-morning and morning hours.”

Noting that other studies have “indirectly suggested that platelets are an important contributing mechanism,” cardiologist Syed M. Jafri says he has recently charted the daily cycle of changing platelet stickiness in nine healthy individuals and three people with chronic chest pain, or angina. He and his co-workers at the Henry Ford Hospital's Heart and Vascular Institute in Detroit tracked blood levels of beta-TG and another natural marker of platelet aggregation, called platelet factor 4.

Jafri says the findings, which he plans to present in Amsterdam this June at the International Congress on Thrombosis and Hemostasis, confirm what earlier studies had suggested: Platelet stickiness reaches a daily low overnight, then begins a steep climb when a person rises. Although reduced blood flow can result from activation of either platelets or a separate blood-coagulation system, Jafri's new data indicate that only platelet stickiness varies with the time of day.

— J. Raloff

Huge black hole may lurk in nearby galaxy

Three years ago, a trio of astronomers began observing a celestial object about 300 million light-years from Earth, the likely product of a collision between two galaxies. They had hoped to study the puzzling source of its brilliant, infrared light. Instead, they stumbled onto a far weightier enigma: evidence hinting at the possible presence of the most massive black hole ever postulated to reside within a galaxy. Black holes are dense, compact objects believed to exist but never definitively detected.

Jonathan Bland-Hawthorn of Rice University in Houston and his two co-workers maintain that the simplest explanation for their observations appears to be a black hole as massive as all the visible stars in the Milky Way, yet compressed into a region just one-ten-thousandth our galaxy's volume. But they agree with other researchers that an unusual feature in the relatively nearby galaxy, named NGC 6240, may have more mundane explanations.

Indeed, François Schweizer of the Carnegie Institution of Washington (D.C.) says, “I think there's only a one in 10 chance there's a black hole there.”

The astronomers embarked on their odyssey using a special instrument, a Fabry-Perot interferometer. Acting like a highly selective filter, it uses the wave-like properties of light and a variable gap between two polished mirrors to pick out whatever visible-light wavelength the astronomers choose to view. Attached to the 2.2-meter University of Hawaii telescope on Mauna Kea, it enabled Bland-Hawthorn and his colleagues to simultaneously chart the velocity of hydrogen-gas atoms throughout much of the telescope's field of view, allowing the first visible-light map of a predominantly infrared-emitting galaxy.

In the April 10 *ASTROPHYSICAL JOURNAL LETTERS*, the scientists report evidence for two rotating disks of gas in this galaxy. One disk orbits around two light-emitting centers at a speed governed by the mass of the ordinary stars and gas within it. In contrast, measurements from another region some 19,000 light-years away show evidence for a second disk with unusual properties.

The team did not directly view the second disk, but deduced its existence from velocity measurements indicating the presence of a rotating body of gas. From its outer to its inner edge, the gaseous disk increases its rotational speed by more than 400 kilometers per second, the researchers found. They also noted a rise in luminosity toward its center, an indication that more hydrogen gas clusters there.

After a colleague confirmed their results last year, the researchers used elementary physics to deduce the gravita-



Keel, P. L. Shoppe

Ellipse marks position of central gaseous disk in NGC 6240. Rotational velocity of a second disk (not shown) suggests the presence of a black hole (cross).

tional tug of an extremely massive, dark and compact object — between 40 billion and 200 billion solar masses — hidden in the region enclosed by the disk.

Cramming several trillion brown-dwarf stars or neutron stars into the tiny region enclosed by the disk would also explain the findings, Bland-Hawthorn says. But he suspects such a concentration of stellar material would not survive without collapsing into a black hole.

William C. Keel has also extensively studied NGC 6240. An astronomer at the University of Alabama in Tuscaloosa, he notes that the character of this galaxy — believed to be in the final throes of forming from the merger of two others — must be carefully considered in interpreting the current work. The black-hole scenario depends on the assumption that Bland-Hawthorn's team really detected a rotating disk of gas, he explains. “If they're being fooled, if the motion in this merging system [merely looks like a disk], then this is just a case of inappropriate interpretation.”

Schweizer says several possible discrepancies point to another explanation. Noting that the researchers admit in their article that parts of the second disk have a relative drop in velocity that can't be explained by simple, planet-like motions, he questions their ability to deduce the presence of a black hole. Instead, Schweizer suggests that material blown out radially from a common center might account for the velocities measured by the researchers. To date, neither Bland-Hawthorn nor co-worker Andrew S. Wilson from the University of Maryland in College Park have ruled out such a possibility.

Bland-Hawthorn told *SCIENCE NEWS* his team hopes to study NGC 6240 with the X-ray satellite ROSAT. Together with higher-resolution observations they made from Mauna Kea last month, these new data might resolve the galactic mystery by the end of the year. — R. Cowen