

Underwater surveys located one such outpost, known as Stingray Lagoon, in 1991. Excavations conducted through this year have yielded numerous well-preserved items once used in salt production, McKillop reports. Inhabitants of Stingray Lagoon apparently boiled seawater in large, thick-walled open bowls, each of which sat on three bolts embedded in a clay base. Numerous examples of all these artifacts emerged from the site, the Louisiana anthropologist says.

Investigators also found abundant pieces of charcoal and the remains of a hearth, clear signs of extensive fire use.

Many ceramic artifacts at Stingray Lagoon, such as pots bearing distinctive stamped designs and figurine whistles, apparently came from inland communities that traded with the salt producers, McKillop holds.

Salt probably left the site in bulk quantities, she adds. The lack of fish bones at Stingray Lagoon indicates residents did not use salt to dry fish for transport elsewhere.

Several additional underwater Maya sites in the vicinity of Stingray Lagoon show signs of less intensive salt production, McKillop says. One of them, known as Wild Cane Cay, has also yielded obsidian objects and other items intended for elite groups. That settlement may have served as one hub for regional trade routes, she suggests.

— B. Bower

Minor climate change can unravel a forest

A modest climate cooling several hundred years ago upset the balance of tree species inhabiting southern Canada, suggesting that even changes spread over several centuries can dramatically alter forests and reduce their productivity, two Canadian scientists report.

Ian D. Campbell of Forestry Canada in Edmonton, Alberta, and John H. McAndrews of the University of Toronto studied how forests in southern Ontario responded to the Little Ice Age, a global cooling that lasted from roughly 1200 A.D. to 1850 A.D. in Canada. Studies of pollen deposited in lakes during that period suggest that the predominantly beech tree forest gave way first to oak and then to pine trees after the start of the cooling. But while pollen can reveal the composition of an ancient forest, it cannot provide information about the land's productivity — the number of trees present.

To test the impact of the climate cooling on woodlands, the researchers simulated the period using a forest model run on a supercomputer. Campbell and McAndrews mimicked the Little Ice Age by gradually lowering the mean annual temperature in southern Ontario by 2°C over the 650-year cool span.

The simulation showed a succession of

Evidence of disks in middleweight stars

For years, stars just a few times more massive than the sun have seemed the neglected stepchildren of astronomers. Deemed too massive to form planets yet too lightweight to strongly influence the interstellar medium, these stars' evolution was largely overlooked by researchers. Now, astronomers have for the first time caught some 2,000 intermediate-mass stars in the act of forming.

Ranging from three to eight times the mass of the sun, these stars belong to a Milky Way cluster known as NGC 6611. Based on visible-light and infrared observations, astronomers find indirect evidence that some 20 percent of these newborns have disks or spherical envelopes of dust surrounding them, says Lynne A. Hillenbrand of the University of Massachusetts at Amherst.

That compares with the 60 percent of low-mass stars thought to have disks (SN: 1/16/93, p.36). But the new observations may have found only the tip of the iceberg. Intermediate-mass stars have a far shorter youth and their disks may last for just a million years — one-tenth the average lifetime of dusty material encircling a low-mass star — Hillenbrand says. In fact, she adds, the study hints that disks play as important a role in the evolution of intermediate-mass stars as they seem to play for less massive ones.

Hillenbrand and her colleagues report their study in the November *ASTRONOMICAL JOURNAL*.

Disks around sun-like stars may survive long enough to provide the raw



Stars in the young cluster NGC 6611, with the Eagle nebula at center.

material for planets, but those around heavier bodies are most likely dragged into the stellar core before planet making begins. Regardless of mass, however, the formation of a disk may funnel material into a developing star and hasten its growth, Hillenbrand speculates.

The team also finds evidence that some members of cluster NGC 6611, known as Be and Ae stars, are much younger than believed. Astronomers had thought that bright emission lines in the spectra of these stars stemmed from mass loss that occurs when the stars reach about 100 million years of age. But that's inconsistent with the youthfulness of the rest of the cluster stars, which are no more than a few million years old.

The researchers suggest that the Ae and Be stars in the cluster are newborns and that starbirth processes account for the spectra.

— R. Cowen

tree types resembling the pattern of change recorded by ancient pollen. Going beyond the pollen record, the computer experiment also provided a measure of the total amount of biomass within the forest. The biomass declined by 30 percent during the Little Ice Age, indicating that the cooling knocked the region's woodlands far out of equilibrium.

"Nobody had ever demonstrated before that such a minor climate change could have such a large impact," Campbell says.

The finding relates to concerns about future global warming, he notes, because it indicates that climate change does not simply alter the mix of trees in a forest by swapping cool-temperature species for warm, or vice versa. Instead, the slow cooling of the Little Ice Age caused a cascade of reactions that is taking centuries to play out.

The forest biomass declined steeply because the disappearance of beech trees cleared large sections. This prompted the spread of oak, which thrives in open spaces. Once the forest thickened, the oak died back, giving way

to more shade-tolerant pine trees. By the end of the simulation, the year 2000, the forest still had not recovered its former productivity, the researchers report in the Nov. 25 *NATURE*.

Climate experts debate how the Earth will react to the buildup of greenhouse gas pollution in the atmosphere, but most predict that the warming will proceed several times faster than the cooling modeled by Campbell and McAndrews.

"Most people are predicting fairly significant and rapid climate change. It's a warming instead of a cooling. But my research shows it was the rapidity of the change that was important, not the direction," Campbell says.

Some ecologists note that Campbell and McAndrews did not demonstrate that the model they used accurately represented how the real forest responded to the Little Ice Age. In particular, the simple model did not incorporate wildfire and other types of disturbance that other researchers have shown to play an important role during shifts in climate.

— R. Monastersky