

tsunami. In this expedition, they mapped the seabed and drilled samples of sediments. In late February, they used a robotic sub to photograph the seafloor.

The scientists described their findings at a July meeting of the International Union of Geodesy and Geophysics in Birmingham, England, and in the July 27 *Eos*.

The exploration focused on the continental slope, which plunges into a 4-kilometer-deep trench. This chasm marks where a piece of the Pacific Ocean floor crashes into New Guinea. As the Pacific tectonic plate scrapes beneath the island, it creates the deep trench and sparks frequent earthquakes.

The survey found that the extremely steep continental slope bears a thick carpet of sediments. In places, this coating has slid downhill in speedy landslides and slower-moving slumps.

On one dive, the researchers discovered a fresh, amphitheater-shaped scar, created when a giant chunk of sediment slumped downhill. "We know this slope failed sometime in the past," says Watts. "The mystery is, When did it go?"

The researchers also found a 15-km-long fault that showed evidence of recent movement. Faulting of the seafloor can generate tsunamis when one side jumps up and the other side drops, displacing water in the process.

Tappin, Watts, and others who think a slump caused the waves contend that the fault could not be the source. Eyewitness accounts indicate that the first wave struck shore about 20 minutes after the main shock of the earthquake, too long for the tsunami to have originated from subsea faulting during the quake. A slump, however, typically lags several minutes after an earthquake and could explain the delay.

When Watts modeled the tsunami, he obtained better results using an undersea slump than a sea-bottom quake. "I'm convinced that the main part of the tsunami was generated by one giant slump," he says.

Further support comes from a 70-second-long rumble recorded in the middle of the Pacific soon after the earthquake. This sound lasted too long to have come

from a small aftershock and may represent a seafloor slide, says Emile A. Okal of Northwestern University in Evanston, Ill.

Some survey participants, however, discount the slump theory. Harry Yeh of the University of Washington in Seattle argues that simulations of a subsea quake can explain the tsunami's size. The team found evidence for small slides but no obvious signs of a giant slump, he says.

Whatever the outcome of the debate, the recent tsunami is forcing researchers to consider slumps as potential sources of giant waves, says Eddie Bernard, coordinator of the National Tsunami Hazard Mitigation Program in Seattle. "I think the good thing that this has done is to open our eyes."

The disaster also suggests that relatively modest quakes, such as the kind that occasionally rock southern California, can trigger giant tsunamis by setting off slides. "This makes the hazard much more dangerous than the scientific community has perceived it in the past," says Bernard.

—R. Monastersky

Hormone mimics: New assessments air

Dishwashing detergents, pesticides, and even contraceptives contain nonylphenols or compounds that break down into nonylphenols. These chemicals are members of a group that mimic hormones and appear to harm wildlife. Nonylphenols taint waterways throughout the world, especially those downstream of municipal waste treatment plants.

Given half a chance, however, waterborne nonylphenols will take to the air, a new study finds. Their evaporation from water allows the chemicals to travel long distances before settling down again—potentially on land far from water, notes study leader Steven J. Eisenreich of Rutgers University in New Brunswick, N.J.

Using a chemical fingerprinting technique known as gas chromatography, his team identified pollutants in water in the Hudson River estuary and in air nearby. Their work, reported in the Aug. 1 *ENVIRONMENTAL SCIENCE & TECHNOLOGY*, turned up both water and air pollutants with "an unequivocal match" to the signature of nonylphenols.

The data, the first showing these pollutants in the atmosphere, detected airborne amounts ranging from just above zero to 70 nanograms per cubic meter, which the researchers regarded as a "high concentration." Eisenreich says, "Water was definitely the source of these chemicals in the air."

The Rutgers chemists suspect airborne nonylphenols are "ubiquitous" worldwide. Because nonylphenol concentrations in some European rivers are 10 to 100 times as high as in the Hudson estuary, the airborne chemicals are per-

haps even more prevalent elsewhere in the world than in the area studied. The air data raise concern about new routes of human exposure, the scientists say.

Their finding disturbs Susan Sang of the World Wildlife Fund Canada in Toronto, which advocates a phaseout of surfactants that degrade to nonylphenols. When nonylphenol concentrations in water diminish, it has looked like the pollutants were breaking down, Sang says. "It now appears they were just evaporating and moving to where you wouldn't have expected to find them," she says.

Fish exposed to nonylphenols have developed reproductive and other abnormalities (SN: 5/8/99, p. 293). Because of such findings, Sang notes, Canadian officials have recommended that pregnant women avoid nonylphenol exposure.

The Canadian government is also assessing nonylphenol risks. If its findings, due next spring, indicate the pollutants are toxic, the government could require monitoring or even limit nonylphenol release, notes Philippa Cureton of Environment Canada in Hull, Quebec.

Determining whether such pollutants pose risks to people, however, will generally require much more research, concludes a panel convened by the National Research Council (NRC) in Washington, D.C. It released a report of its 4-year assessment of the toxicity of hormonelike chemicals last week.

"We couldn't find any clear evidence that people had been harmed by typical environmental exposures to hormonally active chemicals," observes panel mem-

ber James C. Lamb IV, a consulting toxicologist in Reston, Va.

Then again, few studies have probed for effects in humans, argues panelist Ana M. Soto of the Tufts University School of Medicine in Boston. A further limitation of the new assessment, she maintains, was its "focus on correlations between one chemical and an effect." Most people face coincident exposures to several hormone mimics—such as nonylphenols, phthalates, and PCBs—and her own studies indicate that the effects can be "at least additive," she says.

When it comes to wildlife, "there was clearly evidence of very negative reproductive effects in populations exposed to chemicals commonly called endocrine disruptors," notes panelist Joanna Burger from Rutgers. The NRC consensus report "doesn't dismiss these findings," she says. Indeed, she notes, it documents many of the hormonal mechanisms that are likely responsible.

Though many pollutants may exhibit hormonal activity, few studies have proved that such hormonal action is responsible for the toxicity of these agents, says panel chair Ernst Knobil from the University of Texas Medical School at Houston. He says that one of his committee's "most radical" actions was to "abandon the term *endocrine disruptor*" to describe hormone mimics.

To Lamb, this decision indirectly challenges the value of a new federal program to screen all commercial U.S. chemicals for hormonal action (SN: 10/17/98, p. 251). He says, "I'm concerned that we'll spend all this money chasing hormones—with no certainty that it will help us predict risks."

—J. Raloff