(SN: 12/28, p. 639). Clouds of it were seen in the direction of the galactic center by a University of California group including Drs. A. C. Cheung, D. M. Rank, C. H. Townes, D. D. Thornton and W. J. Welch.

Now the same group reports in the Feb. 15 NATURE that while looking for more ammonia, they not only found it but found water as well. The water appeared in three directions: those of the Orion nebula, the object called Sagittarius B2 and the object called W49.

The water, like the ammonia, is identified by its characteristic emissions in the microwave radio spectrum. "The radiation found is attributed to H₂O because its frequency coincides very closely to that found for H₂O in the laboratory, and no other known atomic or molecular species can explain the observations," say the astronomers.

Molecular spectra are complex, how-

Molecular spectra are complex, however, and in the case of Sagittarius B2, where ammonia was found as well as water, the question may arise whether the water-identifying wavelength might not be attributed to the ammonia instead. Dr. Cheung and his associates argue that it would be extremely difficult for the ammonia to be excited in the way necessary for it to produce the disputed wavelength with the observed brightness. Therefore the wavelength is most likely to come from water, they say.

Furthermore, they point out, strong water emission comes from the Orion nebula, where no ammonia is seen.

silicate compounds seem to appear in the region around the star 119 Tauri. The infrared spectrum of this star as it reaches the earth shows absorption at 9.7 and 10.6 microns. These absorptions, say Drs. R. F. Knacke and J. E. Gaustad of the University of California at Berkeley and F. C. Gillett and W. A. Stein of the University of California at San Diego, could be made by a cloud of silicate material, a combination of either iron or magnesium with silicon trioxide.

Dr. Knacke and his associates surmise that such compounds might have been formed inside the star, a cool supergiant in which various compounds are known to exist, and then blown into space by some disturbance.

Dr. Gaustad and Dr. William C. Saslaw suggest that the interstellar dust grains may be diamond crystals. The classical models for the dust, usually ice or ice-coated graphite, do not fit all the modern observations, they say. Carbon in the form of diamond would.

But there is a major objection: Diamond normally forms only under heavy pressure, a condition opposite to those in interstellar space. Under pressureless conditions carbon would be expected to crystallize in other forms.

To deal with the objection, Drs. Gaustad and Saslaw point out that a diamond crystal has many more points at which carbon atoms could stick than graphite does. In the conditions of random collisions in interstellar space, they say, this means that diamond crystals would have a better chance of growing than graphite. Thus diamond might grow under interstellar conditions even if it were thermodynamically unstable and after a certain lifetime changed to some other form.

To determine whether the dust is certainly diamond, say Drs. Gaustad and Saslaw, it will be necessary to make detailed studies of ultraviolet absorption by the dust and of the mechanisms of diamond formation under unusual conditions.

SEALAB

Board of Inquiry

With evidence consisting of a coroner's report and a small empty can, the Navy last week prepared to name a board of inquiry into the death of Sealab III aquanaut Berry L. Cannon (SN: 3/1, p. 210).

In the coroner's examination Cannon's blood was found to contain excess amounts of carbon dioxide. This, Navy doctors say, could have given the diver convulsions in "a matter of minutes" and caused fibrillations of the ventricles of his heart. Additional evidence of CO2 poisoning was the discovery that one of the Sealab diving units had an empty Baralyme air-purification canister. These are supposed to remove carbon dioxide as breathing mixtures are recycled. But because the diving units are not marked for each aquanaut, it will be up to the board of inquiry to determine if the empty canister was indeed Cannon's.

After months of delays due to gas leaks and mechanical problems, the climactic stage of the Sealab program was finally about to get underway, when Cannon's tragic and sudden death on Feb. 17 brought things to an abrupt halt. But for the tragedy, nine-man teams of aquanauts that week would have begun living in 12-day shifts, 620 feet down in the Pacific off San Clemente Island. Last week, however, the habitat and most of its crew were to be returned to the California mainland to wait out the investigation.

Meanwhile, project officials hope to get approval to resume testing of Sealab's experimental diving system, including the elevator-like personnel transfer capsules, decompression chambers on the deck of the surface support ship and the divers' own equipment, but without the habitat, while the inquiry is still in progress.



8 march 1969/vol. 95/science news/235