

Skyscrapers as Lightning Rods

Tall buildings and lightning rods mounted on high towers protect neighboring structures from lightning, provided they are not so high as to extend out of the cone of protection. This protected area extends around the base of the high building for a distance of between two and four times its height. Imaginary lines drawn from the top of the building to the edge of the protected area define the protected cone, says F. W. Peek, Jr., in charge of the General Electric Company's high-voltage investigations.

Mr. Peek's investigations have been made with artificial lightning at pressures of as high as three and a half million volts. These man-made flashes have been used on small models of buildings. However, confirmation of his discoveries was obtained by studying a natural electrical storm that occurred in New York last summer, and during which the New York World building was struck. Though this building is close to the Woolworth Tower and is in the 1,100-foot circle around its base that is pro-

Physics

tected, the dome of the World building extends for about a hundred feet outside the cone, and that is the reason that it was struck, explains Mr. Peek. If it had been 200 feet closer the Woolworth building, it would have been protected.

Practical application of these experiments, says Mr. Peek, has already been made in California, in safeguarding oil storage tanks from lightning. Several tall rods, placed outside the big reservoirs, provide overlapping cones of protection and reduce the danger to a minimum.

Science News-Letter, May 26, 1928

Instruments Record Quake

By means of records obtained on seismograph instruments at observatories as distant as Massachusetts, Alaska, Hawaii and the Philippine Islands, and gathered by Science Service, the earthquake experts of the U. S. Coast and Geodetic Survey here have located the center of the quake on Monday afternoon, May 14. It was at latitude 8.0 degrees south and longitude 80.5 degrees west, and occurred at 5 hours 14 minutes 20 seconds p. m., Eastern Standard Time. This location is in the Pacific Ocean about 70 miles off the Peruvian coast. Fortunately for residents of this region, the slip of the ocean bed which produced it was mostly in a horizontal direction. Had the shaking been up and down, a severe tidal wave would doubtless have been the result.

At Guayaquil, Ecuador, nearly four hundred miles to the north of the center of the shake, severe shaking was reported. This, and the fact that seismograph observatories at such distant points recorded it, shows that it was a quake of great severity.

The seismograph stations that reported their records of the quake to Science Service were those of the U. S. Coast and Geodetic Survey at Tucson, Arizona; Sitka, Alaska; Honolulu, T. H., and Chicago, Ill.; of the Jesuit Seismological Association at Georgetown University, Washington; Fordham University, New York; Regis College, Denver; Loyola University, New Orleans, and St. Louis University, St. Louis, and the stations of Harvard University, Cambridge; the Dominion Observatory, Ottawa, Canada, and the Manila Observatory in the Philippine Islands.

Science News-Letter, May 26, 1928

NATURE RAMBLINGS

BY FRANK THONE

Natural History



III Weeds

Now is the time when schools and churches and any other groups of people who have any excuse at all for getting out of doors plan picnics. And by an evil coincidence poison ivy, about the most universally distributed plant pest in America, is coming into bloom, and according to widespread opinion is at the climax of its virulence.

The plant is not hard to recognize, if you will look about a bit before spreading your picnic cloth or hanging up your hammock. It grows as a low, little branched shrub out of the ground, or climbs up the trunks of trees as a close-clinging vine, with innumerable aerial roots with which to hang onto the bark. In either case, however, its leaves are the same; compound affairs with three broadly ovate leaflets; really rather attractive things, with their glossy green shine. The flowers are slender, irregular pyramids of tiny white bloom.

It is very easy to get poisoned with poison ivy; that happens on the mildest of country walks, for poison ivy is everywhere. Poison sumac is reserved for slightly hardier souls, who go in for hiking or nature-study activities that may require wet feet, because poison sumac is a creature of the bog-edges, and does not grow in upland woods at all. This is perhaps fortunate, for though fewer persons are susceptible to it, the luckless ones it does affect get a much worse "dose", usually, than poison ivy is able to inflict.

Poison sumac is easy enough to identify. First, it is always in low ground, usually in wet ground. Second, it looks much like ordinary sumac, except that its bark is a pallid gray. Third, its fruits (which persist on their stalks for a long time) are quite unlike those of ordinary sumac, being round, white, somewhat waxy (*Turn to next page*).

Ornith evolution

Palaeontology

The Archæopteryx, a primitive bird, is believed to have evolved from a reptile.

Ma Sauropod, a reptile odd,
In ages Carboniferous,
Had made her nest; had done her best;
And then became vociferous.

"Oh, Pa!" cried she, "Do come and see!
I shouldn't set, I warn you!
I hesitate to incubate
The funny eggs I've borne you!"

"Pooh, pooh!" said Pa, "What ails you Ma?
Don't tell me you're afraid
To try and hatch the little batch
Of progeny you've laid!"

"Oh, well," she hissed, "If you insist,
I'll hatch the horrid brood!"
To end the spat she sat and sat,
And hatching soon ensued.

Ma Sauropod exclaimed, "Ecod!
What's this my gaze transfixes!
I feared as much! Oh, what a clutch
Of Archæopteryxes!"

Richard Ashman.

Science News-Letter, May 26, 1928

Bacteriophage as a Specific Antigen

Bacteriology

J. BRONFENBRENNER, in *The Newer Knowledge of Bacteriology and Immunology* (Univ. of Chicago Pr.):

It has just been suggested that the particles seemingly endowed with the power of initiating transmissible lysis may not represent autonomous "corpuscles" of a parasite, but serve merely as a vehicle on which the active principle is absorbed. The active agent itself, however, may still be particulate, i. e., may consist of minute aggregates composed of a number of molecules each, and these small aggregates may still possess all the characteristics previously ascribed to larger carrier-particles. These aggregates, though perhaps considerably smaller than $20 \mu\mu$ in diameter, may still represent, according to the definition of D'Herelle, "colloidal micellae which differ from other forms of life only in their size." That the active agent may not be a crystalloid is suggested, for instance,

III Weeds—Continued



berries in drooping clusters instead of the crowded erect masses of black seeds of the upland sumacs.

It is possible to insure oneself against ivy and sumac poisoning. If you know you are susceptible, wash your hands and face in a five per cent. solution of ferric chloride in water—your druggist can mix it for you—and let it dry on the skin. A little glycerine added will make it more agreeable to apply. That will neutralize the poison when you meet it.

If you have not taken this precaution, and find yourself poisoned, then use a three to five per cent. solution of potassium permanganate, which your druggist can also supply. Swab it on with a bit of cotton. It will sting a bit, but it will kill the poison. The brown stain on the skin can be removed afterward with a one per cent. solution of oxalic acid.

Science News-Letter, May 26, 1928

by its behavior with respect to solubility. Once dried, the preparations of bacteriophage cannot be brought to the state of initial dispersion in water. This is particularly true of purified preparations of bacteriophage deprived of all, or nearly all, of the extraneous material. The colloidal nature of the active agent is further evidenced by the fact that it acts as an antigen and causes the production of specific antibodies when introduced parenterally into experimental animals. While the antisera thus obtained are capable of neutralizing the activity of the bacteriophage, they usually show simultaneously the presence of agglutinins, opsonins, precipitins, and complement-fixing antibodies for homologous bacteria. Moreover, purely antibacterial sera apparently possess the power to inhibit the progress of lysis by homologous bacteriophages. In spite of the inference suggested by these findings, it has been possible to establish that the active principle possesses independent (autonomous) antigenic value. Thus, when sera prepared by immunization with filtrates of lysed cultures are absorbed with homologous bacteria, they no longer react with bacteria and their products, but retain their power to neutralize the bacteriophage unimpaired. If the filtrates of lysed cultures are purified by removal or digestion of bacterial protein, leaving only the active principle, the anti-sera obtained by immunization with such purified preparations do not react with bacteria, while they do neutralize the phage. Moreover, in the case of phages capable of acting on several related species of bacteria, the antigenic specificity of the filtrate is constant for each phage and is independent of the bacterial substratum which was used to obtain the active filtrate.

As for the phage-inhibiting activity of antibacterial sera, it has been demonstrated that this action is not directed against the phage, but against the bacteria, rendering them less susceptible to lysis. It has been suggested that this effect is probably due to the saturation by agglutinins of receptors susceptible to union with phage.

It appears then that when filtrates of lysed cultures are used in immunization of animals, the agent of transmissible lysis acts as a specific antigen, independently of other anti-

gens, notably those of bacterial origin, present in the filtrates at the same time. The question naturally arises: Is the active agent therefore to be considered an autonomous, organized being, totally foreign to bacteria and multiplying at their expense, or can it still be a product of bacteria, accumulating in the culture during the growth or during the lysis of bacteria, but antigenically distinct from them?

Science News-Letter, May 26, 1928

New flying rules established by the Department of Commerce prohibit acrobatic maneuvers at less than 1,500 feet.

The barking sands in the Hawaiian Islands give forth a sound like the barking of a dog when rubbed together or slid over.

The question of whether the exhibition of mummies of ancient kings is disrespectful to their dignity has been raised in Egypt.

BINDER COVERS FOR SCIENCE NEWS-LETTER

Many subscribers have expressed a desire for a convenient binder in which to file their copies of the *Science News-Letter*. We therefore have prepared an attractive and durable loose-leaf binder-cover of gray leather-like stock, printed in dark green and complete with fasteners. Each binder-cover will hold one volume (six months or 26 issues).

To facilitate punching the issues of the *Science News-Letter* to fit this binder-cover, a pattern showing where holes should be placed appears each week on the back cover page.

To obtain a binder-cover, send 20 cents in stamps (make them 2s, please), together with your name and address (please print) to

SCIENCE SERVICE
21st and B Sts.
Washington, D. C.