

METEOROLOGY

Fallout Patterns

► **KNOWING** which way the wind blows is not always helpful. At least not where deadly radioactive fallout is a factor, the Federal Civil Defense Administration warns.

Cautioning the public and civil defense personnel alike, the Administration reported that it is high altitude winds, up to 80,000 feet, that have the greatest effect on the direction of fallout. Surface winds, FCDA scientists point out, are no indication of the course radioactive fallout takes after an atomic or hydrogen bomb attack.

The government agency also warned that the "flat assumption that the fallout pattern is invariably in a west-to-east elliptical pattern" is wrong.

The "typical" west-to-east cigar-shaped pattern can easily be reshaped into a spilled-ink-like mass and blown in any direction, tests have shown. Although prevailing winds in the nation are west-to-east, they can blow the other way, particularly in the summer.

In issuing the warning, the FCDA reports that rain and snowfall would reduce the

area of fallout, but at the same time, increase the concentration of radioactive particles in some areas.

At present, the U. S. Weather Bureau is issuing twice daily forecasts for all critical target areas on fallout direction and arrival time.

The civil defense officials included their warnings in a technical bulletin, "Fallout and Winds." In the bulletin, they list five rules-of-thumb about the subject:

1. Up to 40,000 feet, the prevailing winds are out of the west, particularly in the spring, fall and winter.

2. Strongest winds are usually found at 30,000 to 40,000 feet and these would determine the general direction and length of the pattern.

3. "Typical" wind direction varies more in the summer, especially in southeastern United States.

4. Above 60,000 feet, winds from the east occur frequently in all seasons.

5. Winds blow less constantly from the west along the Pacific Coast.

Science News Letter, January 21, 1956

track's curvature caused by the magnetic field indicates the particle's mass.

Measurement of both energy and mass simultaneously has been virtually impossible because the curved track becomes distorted.

The Berkeley scientists modified the cloud chamber to use helium instead of air. Since helium atoms each have only two electrons to be knocked off by cosmic ray particles, the track of droplets is thinner. No electrical field is needed to separate negatively from positively charged droplets, and the track's distortion is eliminated.

Science News Letter, January 21, 1956

SCIENCE NEWS LETTER

VOL. 69 JANUARY 21, 1956 NO. 3

The Weekly Summary of Current Science, published every Saturday by SCIENCE SERVICE, Inc., 1719 N. St., N. W., Washington 6, D. C., North 7-2255. Edited by WATSON DAVIS.

Subscription rates: 1 yr., \$5.50; 2 yrs., \$10.00; 3 yrs., \$14.50; single copy, 15 cents, more than six months old, 25 cents. No charge for foreign postage.

Change of address: Three weeks notice is required. When ordering a change please state exactly how magazine is now addressed. Your new address should include postal zone number if you have one.

Copyright, 1956, by Science Service, Inc. Reproduction of any portion of SCIENCE NEWS LETTER is strictly prohibited. Newspapers, magazines and other publications are invited to avail themselves of the numerous syndicate services issued by Science Service. Science Service also publishes CHEMISTRY (monthly) and THINGS OF Science (monthly).

Printed in U. S. A. Entered as second class matter at the post office at Washington, D. C., under the act of March 3, 1879. Acceptance for mailing at the special rate of postage provided for by Sec. 34.40, P. L. and R., 1948 Edition, paragraph (d) (act of February 28, 1925; 39 U. S. Code 283) authorized February 28, 1950. Established in mimeographed form March 19, 1922. Title registered as trademark, U. S. and Canadian Patent Offices. Indexed in Reader's Guide to Periodical Literature, Abridged Guide, and the Engineering Index.

Member Audit Bureau of Circulation, Advertising Representatives: Howland and Howland, Inc., 1 E. 54th St., New York 22, Eldorado 5-5666, and 435 N. Michigan Ave., Chicago 11, Superior 7-6048.

SCIENCE SERVICE

The Institution for the Popularization of Science organized 1921 as a non-profit corporation.

Board of Trustees — Nominated by the American Association for the Advancement of Science: Kirtley F. Mather, Harvard University; Paul B. Sears, Yale University; Karl Lark-Horovitz, Purdue University. Nominated by the National Academy of Sciences: Edward U. Condon, Berkeley, Calif.; Harlow Shapley, Harvard College Observatory; George W. Corner, Carnegie Institution of Washington. Nominated by the National Research Council: Ross G. Harrison, Yale University; Leonard Carmichael, Smithsonian Institution; Jerome Hunsaker, Massachusetts Institute of Technology. Nominated by the Journalistic Profession: Neil H. Swanson, Ruxton, Md.; O. W. Riegel, Washington and Lee University; Michael A. Gorman, Flint Journal. Nominated by the Scripps Estate; Edward J. Meeman, Memphis Press-Scimitar; John T. O'Rourke, Washington Daily News; Charles E. Scripps, Cincinnati, Ohio.

Officers — President: Leonard Carmichael; Vice President and Chairman of Executive Committee: Charles E. Scripps; Treasurer: O. W. Riegel; Secretary: Watson Davis.

Staff — Director: Watson Davis. Writers: Jane Stafford, Marjorie Van de Water, Ann Ewing, Horace Loffin, Howard Simons, Dorothy Schriver, Helen M. Davis. Science Clubs of America: Joseph H. Kraus, Margaret E. Patterson. Photography: Fremont Davis. Production: Priscilla Howe, Marcia Nelson. Interlingua Division in New York: Alexander Gode, 80 E. 11th St., GRamercy 3-5410.

PSYCHOLOGY

Young Scientists Studied

► **THE** talented young scientists of America are characterized by capacity, interest and freedom.

This is indicated by a survey of 600 who have won recent honors in the National Science Talent Search, conducted by Dr. Robert D. MacCurdy of the University of Florida, and reported to the American Association for the Advancement of Science in Atlanta.

In personality the outstanding young scientists show leadership, self-control, self-discipline, curiosity, persistence, and complexity of outlook, Dr. MacCurdy's study, done at Boston University, showed.

He found that they recognized the interdependence of science and society. They made their decisions to be scientists very early when they were in elementary school and they show their love of humanity through their participation in science.

Dr. MacCurdy's results were based on inquiries to two years of National Science Talent Search honor winners totaling 600. Based on the experience of earlier Science Talent Search winners over 15 years, 93% of them will become scientists and engineers.

"Scientific manpower is the product of a long process of educational experience," Dr. MacCurdy said. "Career choices are made at a very early age and the peak of productive efficiency is under 35 years of age. Scientists must be identified when very young and provided with an environ-

ment that will encourage science development in a few years."

Youngsters who seem to have promise as scientists-to-be show high drive and a tendency to introversion, Paul Brandwein of Forest Hills High School, N. Y., told the scientists. There are discernible periods of incubation and illumination which seem to be characteristic of the way these young people work.

The teacher who is successful in working with young scientists, aside from his superior training and wide interests, also serves as a father or mother image, he has found.

Science News Letter, January 21, 1956

PHYSICS

Slimmer Particle Tracks In Cloud Chambers

► **RELIABLE MEASUREMENTS** can be made of both the mass and energy of cosmic ray particles slicing through a cloud chamber for the first time by means of a method developed by Drs. William B. Fretter and E. W. Friesen, University of California physicists.

High-energy charged particles passing through cloud chambers leave tracks formed from droplets of condensed air.

Scientists count the droplets to determine the velocity, or energy, of the particles, using a magnetic field to separate the positive from negatively charged droplets. The