


**EYE MIRROR Magnifies 5X**

for "Iri-diagnosis"  
Permits direct viewing of the capillaries in your own eyes or for treating cysts. Greatly helps remove dust from the eye or unwanted hair from the area, also for eye make-up. Front-surfaced mirror of finest optical glass polished to highest perfection. Aluminized. Brilliant reflectivity. Useful, beautiful. Diameter 1 1/2" (40 mm). Packaged in lucite. **\$2.50**

**HARRY ROSS** Scientific & Lab Apparatus  
63 Reade St., N. Y. 7, N. Y.

**URNS YOUR HOME WIRING INTO  
A GIANT TV ANTENNA!  
PLUG-IN ANTENNA**



● Plugs in Like an Electric Cord—Uses No Electricity

● Eliminates "Rabbit Ears" and Outdoor Antennas --Stays Out-of-Sight

● Gives Clear, Sharp Reception

**ONLY... \$2.98**  
Reg. \$3.98 **2**ppd.

Attach one end to TV set and plug other end into electrical outlet and INSTANTLY you'll

enjoy sharp, clear pictures. Replaces old-fashioned ugly indoor and outdoor antennas. Uses no current, never interferes with use of appliances, never wears out or needs replacement.

Satisfaction is Guaranteed


**DAMAR'S** 75-D Damar Bldg. Elizabeth, N. J.

**HERE'S A QUALITY  
STUDENT MICROSCOPE  
AT A BUDGET PRICE!**

Although budget priced, the UNITRON Model MUS is definitely not just another student microscope. It includes these significant features often lacking in much more costly student models:

- NOT JUST a disk diaphragm... but an iris diaphragm for perfect control of aperture.
- NOT JUST a single focusing control... but both coarse and fine.
- NOT JUST a mirror... but a condenser for optimum illumination and resolution.
- NOT JUST one objective... but three: 5X, 10X, 40X.
- NOT JUST one eyepiece... but a choice of two from 5X, 10X, 15X.

Durable, sturdy — ideally suited to withstand the use and abuse of classroom and laboratory.



Only **\$74**

F. O. B. Destination

ASK FOR A FREE 10 DAY TRIAL

Quantity discounts start at 3 — Only \$65.12 each for 25

**UNITRON**

INSTRUMENT DIVISION of UNITED SCIENTIFIC CO.  
204-206 MILK STREET • BOSTON 9, MASS.

Please rush UNITRON's Microscope Catalog. 5B-1

Name \_\_\_\_\_  
Company \_\_\_\_\_  
Address \_\_\_\_\_  
City \_\_\_\_\_ State \_\_\_\_\_

## ASTRONOMY

# Twinkling Stars Studied

ASTRONOMERS at the Mt. Palomar Observatory, Pasadena, Calif., have completed studies of twinkling stars which will help measure astronomical distances far more accurately than ever before.

For more than 40 years astronomers have based their measurements on the twinklers, pulsating stars known as cepheid variables. But the calibrations were not as accurate.

These stars have two unique habits that make it possible to use them as distance indicators. They dim and brighten in a rhythmic cycle, called a period, and the length of their blink period and their brightness are related. In other words, the longer the period, the brighter the star. Twinklers with half-day periods are 100 times brighter than the sun and those with 40-day blinks are 6,000 times brighter than the sun.

This means that if two cepheids of the same period are compared and one of them appears to be brighter than the other, the difference in brightness will be due entirely to the fact that one is farther away than the other. If one appears to be one-fourth as bright as the other, it means that it is twice as far away because brightness varies inversely with the square of the distance. If in addition the actual distance of one of them is known, the distance of the other can easily be calculated.

Astronomers found that a few cepheids twinkled irregularly and that others did not conform exactly to the period-luminosity relationship. Small differences in color were discovered in cepheids having the same period.

A group of astronomers from Mt. Wilson and Palomar Observatories found that some of the cepheids were redder than

others, due to thin dust clouds somewhere between the viewing telescope and the star itself. They also discovered that over a period of many years, a pulsating star loses some of its blueness and becomes yellower and cools off about 1,000 degrees Fahrenheit.

The studies indicate that the larger stars are more reliable as distance indicators. By correcting for the color differences, astronomers can now calculate distances as far as the telescope can see.

Science News Letter, April 2, 1960

## SCIENCE NEWS LETTER

VOL. 77 APRIL 2, 1960 NO. 15

Edited by WATSON DAVIS

The Weekly Summary of Current Science, published every Saturday by SCIENCE SERVICE, Inc., 1719 N St., N.W., Washington 6, D. C., North 7-2255. Cable Address: SCIENSERV.

Subscription rates: 1 yr., \$5.50; 2 yrs., \$10.00; 3 yrs., \$14.50; ten or more copies in one package to one address, 7 1/2 cents per copy per week; single copy, 15 cents, more than six months old, 25 cents. No charge for foreign postage.

Change of address: Three weeks notice is required. When ordering a change please state exactly how magazine is now addressed. Your new address should include postal zone number if you have one.

Copyright © 1960 by Science Service, Inc. Reproduction of any portion of SCIENCE NEWS LETTER is strictly prohibited. Newspapers, magazines and other publications are invited to avail themselves of the numerous syndicated services issued by Science Service. Science Service also publishes CHEMISTRY (eight times a year) and THINGS of Science (monthly).

Printed in U.S.A. Second class postage paid at Washington, D. C. Established in mimeograph form March 13, 1922. Title registered as trademark, U. S. and Canadian Patent Offices. Indexed in Reader's Guide to Periodical Literature, Abridged Guide, and the Engineering Index. Member Audit Bureau of Circulation.



### SCIENCE SERVICE

The Institution for the Popularization of Science organized 1921 as a non-profit corporation.

Board of Trustees—Nominated by the American Association for the Advancement of Science: William W. Rubey, U. S. Geological Survey; Wallace R. Brode, National Bureau of Standards; Douglas Whitaker, Rockefeller Institute for Medical Research. Nominated by the National Academy of Sciences: Harlow Shapley, Harvard College Observatory; Philip Bard, Johns Hopkins University; Henry Allen Moe, John Simon Guggenheim Memorial Foundation. Nominated by the National Research Council: Leonard Carmichael, Smithsonian Institution; John R. Dunning, Columbia University; Benjamin H. Willier, Johns Hopkins University. Nominated by the Journalistic Profession: Michael J. O'Gaden, Providence Journal-Bulletin; O. W. Riegel, Washington and Lee University. Lee Hills, Detroit Free Press. Nominated by the Scripps Estate: Edward J. Meeman, Memphis Press-Scimitar; Frank Ford, Washington, D. C.; Charles E. Scripps, Cincinnati, Ohio.

Officers—President: Leonard Carmichael; Vice President and Chairman of Executive Committee: Charles E. Scripps; Treasurer: Wallace R. Brode; Secretary: Watson Davis.

Staff—Director: Watson Davis. Writers: Helen Buechl, Ann Ewing, W. T. M. Grigg, Richard Littell, Allen Long, Jane Marye, Tove Neville, Gloria Tefft, Marjorie Van de Water. Science Youth Division: Joseph H. Kraus, Dorothy Schriver, Shirley Moore. Photography: Fremont Davis. Production: Priscilla Howe, Marcia Nelson. Syndicate Sales: Hallie Jenkins. Interlingua Division in New York: Alexander Gode, 80 E. 11th St., GRamercy 3-3410. Advertising Manager: Fred A. Moulton, METropolitan 8-2562.

## CHEMISTRY

## Silicon-Boron Compound Resists Heat, Oxidation

LARGE QUANTITIES of the little known compound tetraboron silicide, B<sub>4</sub>Si, can now be prepared for the first time.

Dr. Ervin Colton of the Allis-Chalmers Manufacturing Company, Milwaukee, Wis., reports in the Journal of the American Chemical Society, 82:1002, 1960, that this compound in reasonably pure form is obtained when a mixture of the elements boron and silicon is heated in an inert atmosphere at between 2,200 degrees and 2,500 degrees Fahrenheit.

Objects made from the powdered compound have been shown to be highly resistant to oxidation at 2,500 degrees Fahrenheit due to a protective film that forms on exposure to the air at high temperatures. It also showed excellent thermal shock resistance in that no cracks appeared when the objects were cooled rapidly from that high temperature to room temperatures several times.

Science News Letter, April 2, 1960