stroy its ability to provoke the formation of disease-preventing "anti-bodies" in the blood of the animal or person to be protected. Crystal violet appears to be a much more advantageous material to use for this purpose than any of the three older chemicals.

One of the disadvantages of preventive treatments now in use for hog cholera has been the necessity of using unmodified disease-bearing serum along with them and hog cholera virus, even under veterinary precautions, is not very welcome on a farm. The new treatment promises to do away with this part of the treatment.

One thing that must be determined,

however, before the crystal-violet vaccine is put into use, is its time-lag. At present it appears to take about two weeks for it to become effective; so its use in emergencies might be under some handicap.

Its cost, for large-scale use, cannot be determined until mass production is undertaken, but it is believed that it will be materially less expensive than the preventive treatments now in vogue.

Crystal violet is a dyestuff belonging to the large anilin family of chemicals. Its common name is used as a matter of convenience instead of its longer technical title: hexa-methyl-para-rosanilin.

Science News Letter, February 1, 1936

## 1935 Wetter Than 1934, Also Somewhat Cooler

**D**ROUGHT and heat were rampant in 1934; broken in 1935. Popular impressions on these points receive confirmation and emphasis from a comparative study of weather records of the two years, which has just been completed by the U. S. Weather Bureau.

For the year as a whole, the meteorologists state, the temperature averaged somewhat higher than normal all over the country, except for two limited areas of slightly below-normal temperatures in the Northeast and Northwest. Yet they were still below the quite abnormally high figures for 1934.

The Bureau scientists summarize highs and lows: "The lowest temperature reported in 1935 was 51 degrees below zero at Pine River Dam, Minn., but Grantsburg, Wis., was a close second with fifty degrees below, both occurring in January. The highest for the year was 123 degrees at Cow Creek, Calif.,

## • RADIO

February 4, 4:30 p. m., E.S.T.

A FUTURE FOR AMERICA'S BIRDS
AND BEASTS — Ira N. Gabrielson,
Chief of the U. S. Bureau of Biological
Survey.

February 11, 4:30 p. m., E.S.T.
THE CRIMINAL MIND—Dr. John E.
Lind, St. Elizabeth's Hospital, Washington, D. C.

In the Science Service series of radio discussions led by Watson Davis, Director, over the Columbia Broadcasting System.

in July. This compares with a low of 52 degrees below zero at Stillwater Reservoir, N. Y., and a high of 125 degrees in Death Valley, Calif., for the preceding year. The highest and lowest all-time temperature records for the United States are 134 degrees in Death Valley, Calif., and minus 66 degrees in Yellowstone Park, Wyo."

Rainfall for 1935 far exceeded that for the previous year, even in regions where it still fell short of normal. It did fall below normal in two regions usually accounted rather moist, the Great Lakes region and the Southeast. Elsewhere, especially in the greater part of the traditionally drier West, there was rainfall ranging from slightly below to decidedly above normal.

The Weather Bureau study summarizes: "Except very locally, the only section of the country having less precipitation in 1935 than in 1934 was the Pacific Northwest, while on the other hand, large areas, especially the interior valleys and Great Plains, seriously dry in 1934, had approximately normal to considerably above normal precipitation in 1935.

"An outstanding feature of the year's weather was the persistent spring wetness in the interior valleys. At the beginning of 1936, the soil-moisture situation generally was satisfactory over the eastern half of the country, but most of the western half was still needing precipitation, though heavy rains and snows occurred over the Pacific Northwest during the first two weeks in 1936."

Science News Letter, February 1, 1936

GEOPHYSICS

## North America Being Pulled Into Pacific Ocean

ORTH America is being pulled into the Pacific Ocean by the attraction of the dense layers of the earth's crust which are below the bottom of the Pacific. This in substance is the conclusion of an investigation reported by Dr. Ross Gunn of the U. S. Naval Research Laboratory.

The movement results in a great thickening of light material on the forward edge of the moving continent, which well accounts for the growth of mountain ranges like the Sierra Nevada in the Pacific coast states, adds Dr. Gunn in his report to the *Physical Review*.

Measurements on the velocity of sound waves, states Dr. Gunn, indicate that layers under the Pacific Ocean have a density greater than similar layers underlying the continents. This dense mass of material produces a component of gravitational force at an angle to the normal vertical pull of gravity.

It is this sidewise or tangential pull of gravity which is tugging North America gradually westward, according to Dr. Gunn's findings.

The geophysical evidence, Dr. Gunn maintains, substantiates his earlier papers concerning the origin of the solar system.

Dr. Gunn has shown that a star can acquire sufficient angular velocity to split into two parts. As the parts separate, while revolving around one another, enormous tidal forces are set up which cause both parts of the original star to lose pieces of themselves that eventually become planets. These planets, as they fly off, are hotter on one side than on the other.

It is this difference of temperature, Dr. Gunn has maintained, which ultimately accounts for the uneven distribution of the continental and oceanic hemispheres of the earth, the lopsidedness of the earth's magnetic field and the uneven distribution of the density in the various layers of the now-cooled earth's crust.

It is the last point, the much greater density of the earth's crust beneath the Pacific Ocean, that accounts for the tangential gravitational force that moves North America westward.

Tangential forces in regions of sedimentation and crustal weakness, explains Dr. Gunn, cause overthrusting of the outer layers. The overthrusting ultimately results in the observed mountain changes.

Science News Letter, February 1, 1936